



PAPER 83-21

**LATE QUATERNARY MORPHOLOGIC DEVELOPMENT
AND SEDIMENTATION, CENTRAL BRITISH
COLUMBIA CONTINENTAL SHELF**

**J.L. LUTERNAUER
J.W. MURRAY**

1983

© Minister of Supply and Services Canada 1983

Available in Canada through

authorized bookstore agents
and other bookstores

or by mail from

Canadian Government Publishing Centre
Supply and Services Canada
Ottawa, Ontario, Canada K1A 0S9

and from

Geological Survey of Canada
601 Booth Street
Ottawa, Canada K1A 0E8

A deposit copy of this publication is also available
for reference in public libraries across Canada

Cat. No. M44-83/21E Canada: \$4.00
ISBN 0-660-11453-4 Other countries: \$4.80

Price subject to change without notice

Critical readers

**C.F.M. Lewis
G.J. Yereth**

**Original manuscript submitted: 1982-10-1
Approved for publication: 1983-04-15**

CONTENTS

1	Abstract/Résumé
1	Introduction
2	Regional Setting and Morphology of Queen Charlotte Sound
2	Physiographic setting and morphologic description of the Sound
2	Geological and geophysical setting
4	Glaciation
4	Oceanography
5	Sediment sources
5	Field and Laboratory Procedures
5	Field
5	Laboratory
6	Results
6	Grain-size distribution
6	Bottom photographs
10	Sonograms
11	Clay-size sediment mineralogy
11	Heavy minerals
11	Glauconite
14	Skeletal carbonate
14	Noncarbonate organic carbon
15	Gravity cores
15	Continuous seismic profiles
22	Discussion
22	Structural/stratigraphic influence on shelf physiography
22	Ice contact and proglacial features
22	Formation of troughs and banks
23	Extent of ice grounding during the Fraser glaciation
23	Impact of lower sea levels
23	Present sediment dispersal and deposition
25	Environmental and economic considerations
25	Conclusions
26	Acknowledgments
26	References
32	Appendix
	Figures
2	1. Index map
2	2. Major physiographic units in the vicinity of Queen Charlotte Sound. Bathymetric contours in metres
3	3. Bathymetry of Queen Charlotte Sound
3	4. True scale cross-sections of Mitchell's Trough and Finlayson channel
4	5. Maximum bottom current velocities recorded while spring tides were running in June-July, 1977
6	6. Location of grab and core samples and camera stations
7	7. Mean size - skewness - sorting categories used to map sediment distribution displayed in Figure 8
7	8. Sediment distribution in Queen Charlotte Sound
7	9. Plot of the ratio of weight per cent mud to weight per cent sand vs sample depth
8	10. a. b. Camera station No. 437; Cook Bank (40 m) c. Camera station No. 267; Goose Island Bank (37 m) d. Camera station No. 311; Goose Island Bank (50 m)
9	11. a, b, c. Camera station No. 430; Goose Island Bank (35-40 m) d. Camera station No. 435; Cook Bank (73 m)
10	12. a, b. Camera station No. 391; Middle Bank (130 m) c. Camera station No. 424; Goose Island Trough (183 m) d. Camera station No. 427; Goose Island Trough (250 m)
11	13. Sonogram and Deep-Tow seismic records of shallowest part of Goose Island Bank
12	14. Sonogram of central Intertrough Area displaying relict ice furrows
12	15. X-ray diffractograms illustrating representative responses of fractions of bottom samples finer than 2 μ m to various analytical treatments (Cu K α radiation)
13	16. Glauconite pellet and heavy mineral distribution

- 13 17. The photo at left displays a variety of semi-enclosed and test-free glauconite particles found in Queen Charlotte Sound sediments. The tests and pellets range in size from 0.354 to 1.00 mm. On the right are two samples of what are identified as test-free glauconite pellets ranging in size from 0.354 to 0.500 mm
- 14 18. Distribution of skeletal carbonate and noncarbonate organic carbon concentration as weight per cent of total mud plus sand fraction
- 14 19. Vertical distribution of grain size, colour and organic content in cores obtained in Queen Charlotte Sound
- 15 20. Photographs of representative sections of cores obtained in Queen Charlotte Sound troughs
- 16 21. Representative X-ray images of sections of cores obtained in Queen Charlotte Sound
- 16 22. Index map of locations of seismic profile tracklines and side viewing sonar tracklines
- 17 23. Air-gun seismic profile A-A' across central Mitchell's Trough
- 18 24. Air-gun seismic profile B-B' along axis of outer Mitchell's Trough
- 18 25. Air-gun seismic profile C-C' at the bend in Mitchell's Trough
- 19 26. Air-gun seismic profile D-D' across central Mitchell's Trough flats
- 19 27. Air-gun seismic profile E-E' along axis of Goose Island Trough north of Sea Otter Shoals
- 20 28. Air-gun seismic profile F-F' (and deep-tow high resolution seismic profile G-G') across outer Moresby Trough
- 20 29. Deep-Tow high resolution seismic profile H-H' across the northeastern margin of Goose Island Bank
- 21 30. Deep-Tow high resolution seismic profile I-I' across the southern flank of outer Goose Island Trough
- 21 31. Deep-Tow seismic profile J-J' across the northern flank of outer Goose Island Trough

LATE QUATERNARY MORPHOLOGIC DEVELOPMENT AND SEDIMENTATION - CENTRAL BRITISH COLUMBIA CONTINENTAL SHELF

Abstract

The physiography of Queen Charlotte Sound is dominated by banks capped with sand and gravel and troughs floored with gravel, sand and mud. Grounded ice, probably including that of the Fraser Glaciation, sculpted most morphologic features and strongly influenced the present sediment distribution. The oceanographic regime which prevailed when sea level was lower contributed to the formation of valleys at bank margins and the concentration of heavy minerals on banks and probably determined the depth of the present shelf break.

Inlets on the southern mainland coast of the shelf are the principal sources of sediment presently supplied to the shelf. This material, composed of olive green organic- and smectite-rich muds, is accumulating on the shelf almost exclusively in a section of trough adjacent to the mouths of these inlets.

Probably the most mobile terrigenous sediment on banks, fine to very fine sand, is being reworked from glaciogenic deposits and swept onto bank margins. Although foraminifers and molluscan skeletal carbonate in places are dominant components of bank sediments, their concentration across the shelf is highly variable because of differences in wave and current climate, dilution by terrigenous detritus and, possibly, proximity to upwelling, nutrient-rich, slope waters.

The best sorted, nonencrusted finer gravels and clean sands best suited for construction materials are most concentrated on the eastern, shallower part of Goose Island Bank. The most extensive skeletal carbonate beds, a potential source of lime for use in the manufacture of cement, extend for several tens of kilometres in a 10-15 km wide band adjacent to northern Vancouver Island and the Scott Island chain. The highest concentration of heavy mineral rich "black sands" are found at or near the 100 m depth on northern Cook Bank.

Résumé

La physiographie du détroit de la Reine-Charlotte est dominée par des bancs couronnés de sable et de gravier et des creux tapissés de gravier, de sable et de boue. La glace échouée, y compris probablement celle de la glaciation de Fraser, a sculpté la plupart des éléments morphologiques et a fortement influencé la répartition actuelle des sédiments. Le régime océanographique qui dominait lorsque le niveau de la mer était plus bas a contribué à la formation de vallées aux marges des bancs et à la concentration de minéraux lourds sur les bancs; il aurait également déterminé la profondeur de l'accroissement actuel.

Les baies du littoral continental sud du plateau sont la source principale des sédiments qui s'accumulent présentement sur le plateau. Composés de boues vert-olive riches en matières organiques et en smectite, ce matériel s'accumule presque exclusivement dans une section de creux adjacente aux embouchures des baies.

Les sédiments terrigènes les plus mobiles des bancs, soit les sables fins à très fins, proviennent de dépôts glaciaires et sont transportés jusqu'aux marges des bancs. Par endroits, les carbonates formés de foraminifères et de squelettes de mollusques sont les éléments dominants des sédiments des bancs; toutefois, leur concentration varie beaucoup en travers du plateau en raison de la différence dans le régime des vagues et des courants, de la dilution par des matériaux détritiques terrigènes et peut-être de la proximité des eaux ascendantes du talus, riches en éléments nutritifs.

Les graviers et les sables propres non incrustés les mieux triés et les plus utiles pour la construction sont concentrés sur la partie est moins profonde du banc de l'île Goose. Les plus vastes couches de carbonate squelettique, source possible de chaux pour la fabrication du ciment, recouvrent plusieurs dizaines de kilomètres dans une bande de 10 à 15 km de large, adjacente à la partie nord de l'île Vancouver et de la chaîne des îles Scott. La plus forte concentration de "sables noirs" riches en minéraux lourds se trouve plus ou moins à 100 m de profondeur sur la partie nord du banc Cook.

INTRODUCTION

This report presents the results of the first detailed study of the morphology and sediments of a part of the British Columbia continental shelf adjacent to the strongly glaciated mainland coast (Holland, 1964). Earlier sedimentologic studies have been carried out in the area encompassed by Queen Charlotte Sound (Fig. 1), by the University of British Columbia Institute of Oceanography (1963), Murray and Mackintosh (1968), Wiese (1969), and Shell Canada Limited (in the course of exploratory surveys during the late 1960s).

This report, which includes some data presented in an earlier unpublished study (Luternauer, 1972), describes the following aspects of Queen Charlotte Sound sediments: grain size, clay mineralogy, colour and the total concentration of (a) heavy minerals, (b) skeletal carbonate, (c) noncarbonate organic carbon and (d) coarse (0.354 to 0.500 mm) "glauconite" pellets in the sand plus mud fraction. Subbottom character of the Sound is interpreted from short cores and continuous seismic profiles. Features on the seafloor are described from side-scan sonar records and underwater photographs.



Figure 1. Index map.

The laboratory and field data and the available information on the morphology and regional setting of the shelf, form the basis for an evaluation of the influence of (a) late Pleistocene glaciation, (b) structural relationships and faulting, (c) sea level changes, (d) oceanographic climate, (e) present day river discharge and (f) organic productivity on the physiography and/or sediment distribution.

REGIONAL SETTING AND MORPHOLOGY OF QUEEN CHARLOTTE SOUND

Physiographic Setting and Morphologic Description of the Sound

According to Holland's (1964) classification of British Columbia physiography, Queen Charlotte Sound lies within the southern part of the Hecate Depression (Fig. 2) which extends from northern Vancouver Island to southeastern Alaska. This depression is bounded on the east by the Coast Mountains and on the west by the mountains of the Queen Charlotte Islands and the open Pacific Ocean. Major subdivisions adjacent to Queen Charlotte Sound are the Nahwitti Lowland of northern Vancouver Island and Hecate Lowland of the British Columbia mainland. These lowlands are generally below 610 m in elevation and are considered by Holland to be remnants of Tertiary erosion surfaces. Milbanke Strandflat, a rocky platform, largely below 33 m in elevation, lies along the western edge of Hecate Lowland. Culbert (1971) suggested that the strandflat is a "mature erosion surface modified by wave action." Back-cutting erosion by coastal piedmont glaciers, sea level fluctuations coupled with wave abrasion, and frost weathering have been considered instrumental in the formation of similar features in Norway (Holteidahl, 1960, 1970).

The bathymetry and physiographic subdivisions of Queen Charlotte Sound are shown on Figure 3. Geographic names are those used on Canadian Hydrographic Service charts; those enclosed by quotation marks are unofficial. The principal physiographic features in Queen Charlotte Sound are broad banks and troughs. Secondary features include valleys at bank margins and broad channels on bank tops, flats and ridges in troughs, terraces, and isolated depressions. Although some lineations in the Sound match those on the adjacent mainland, the physiography of the shelf and mainland is generally markedly dissimilar in that the floor of the Sound is not dissected by as dense a network of narrow depressions and has a considerably more subdued relief (Fig. 4).

Peacock (1935) considered that the great depth of fiords such as those on the mainland adjacent to the Sound resulted not only from glacial excavation and possible incomplete postglacial rebound but also deep fluvial erosion during a Tertiary uplift. Culbert (1971) claimed local relief on the mainland can more readily be attributed to "block subsidence of the fiord zone."

Geological and Geophysical Setting

Queen Charlotte Sound is adjacent to the modern triple junction of the North American, Pacific and Juan de Fuca plates at the base of the continental slope (Keen and Hyndman, 1979). Geophysical evidence suggests that oceanic crust is migrating to the northwest parallel to the base of the slope off the Queen Charlotte Islands and easterly to south-easterly towards and beneath Vancouver Island (Riddiough, 1977).

Although the Sound is adjacent to the most seismically active areas in Canada (Milne et al., 1978), during the period 1899-1977 no earthquakes of magnitude 5 or greater (Richter scale) have occurred within the Sound (Rogers, 1982).

Rocks on the mainland adjacent to the Sound are mainly Mesozoic and Tertiary plutons (Roddick et al., 1966; Baer, 1967). The dominant structural grain is northwesterly. Small shear zones, commonly parallel to this trend, are abundant, but faults with proven displacement are scarce (Roddick et al., 1966). A major, albeit inactive, dislocation zone apparently extends for approximately 300 km along the mainland coast northwest of Price Island (Fig. 3; Souther, 1966). The orientation of many of the fiords on the mainland coincides with a subordinate northeasterly to easterly grain which may have been established prior to the northwesterly structural grain (Roddick et al., 1966).

Within the Insular Belt (Muller, 1977), which comprises Vancouver Island and Queen Charlotte Islands, the dominant structural grain has been established by northwesterly trending dextral faults, associated folds, and zones of intermediate plutonic rocks which dislocate sedimentary and volcanic rock strata of late Paleozoic to Tertiary age (Sutherland Brown, 1966, 1968; Muller et al., 1974; Roddick et al., 1976).




Figure 2. Major physiographic units in the vicinity of Queen Charlotte Sound. (after Holland, 1964). Bathymetric contours in metres.



Figure 3. Bathymetry of Queen Charlotte Sound. Contoured principally from Canadian Hydrographic Service field sheets compiled during 1958-59 employing Decca-Hi Fix navigation system. Areas 20 km from the coast were compiled from Canadian Hydrographic Service charts and Shell Canada Ltd. bathymetric map for area. Arrow heads indicate locations of bathymetric profiles in Figure 4.




Figure 4

True scale cross-sections of Mitchell's Trough and Finlayson channel. Vertical scale in metres. Refer to Figure 3 for cross-section locations.

Interpretations of the origin of Queen Charlotte Sound have been presented by Chase et al. (1973), Stacey (1973) and Yorath and Chase (1981). The most recent of these suggests that as the area now occupied by the Sound drifted over the Anahim mantle plume (Bevier et al., 1979) during the Tertiary, what may have been a continuous landmass rifted apart to form a basin into which mid and late Tertiary marine clastics were deposited (Shouldice, 1971, 1973).

Glaciation

Beer (1967) estimated that Pleistocene glacier ice reached a height of at least 2300 m on the mainland adjacent to the Sound where some of the largest icefields in British Columbia are found. Substantial late Pleistocene ice thicknesses on the mainland are indicated by the local glacio-isostatic depression of the crust: (a) Armstrong (1966) determined that "land in Kitimat-Terrace valley [on the mainland approximately 200 km north of the Sound] was depressed at least 300 m below its present elevation..." and (b) Andrews and Retherford (1978) reported that the late Pleistocene marine limit along the mainland coast of the Sound is "at least 40 m on Calvert Island and most likely 120+ m."

Howes (1981) established that the maximum observed level of the marine limit achieved immediately following the retreat (12 000–13 000 years ago) of the Fraser Glaciation ranges from a possible 175 m along the northeast coast of Vancouver Island to approximately 20 m on the northwest coast. This relationship led him to suggest that ice "was thicker on the east coast versus the west coast..."

Sutherland Brown (1968) suggested that locally generated late Wisconsin glaciers covered Queen Charlotte Islands and coalesced with the Cordilleran ice sheet somewhere in the western Hecate Depression. He further suggested that ice (and any sediment it might have been carrying) from only the southernmost quarter of the Queen Charlotte Islands was directed into Queen Charlotte Sound by the part of the mainland ice sheet that flowed through the Sound (this contention is supported by Wiese's (1969) preliminary study of the heavy mineral distribution in the northern Sound).

Clague et al. (1982) established that the late Quaternary sea level history of mainland British Columbia and eastern Vancouver Island contrasts sharply with that of the Queen Charlotte Islands and western Vancouver Island. In the former areas relative sea level is known to have fallen from about 13 000 to about 6000–9000 years ago (when in some areas it was 12 m or more lower than at present). It then generally rose through late Holocene time. In contrast, relative sea level on the Queen Charlotte Islands was below the present level from sometime before 13 700 years ago until approximately 9500–10 000 years ago. At the close of the Pleistocene relative sea level then rose until about 7500–8500 years ago when it was generally about 15 m above present. The succeeding drop in sea level (evident also on the west coast of Vancouver Island) apparently occurred in the last 5000–6000 years. The divergence of sea level patterns has been attributed to the collapse of a subcrustal forebulge during ice retreat and tectonic uplift of the outer coast (Clague et al., 1982; Riddihough, 1982).

The following evidence points to a limited and (or) short-lived extension of late Wisconsin ice across the Queen Charlotte Sound shelf and suggests that the outer part of the shelf may have undergone only minor glacio-isostatic depression during the Fraser Glaciation: (a) northern and southern mainland late Wisconsin coastal glaciers achieved their maximum extension about 15 000 years B.P. but had retreated from the outermost mainland coast (but not necessarily the fiords) by about 12 000–13 000 years B.P. (Clague et al., 1980; Clague, personal communication, 1983);




Figure 5. Maximum bottom current velocities recorded while spring tides were running in June–July, 1977. Extracted from data supplied by R.E. Thomson (Institute of Ocean Sciences, Sidney, B.C.).

(b) northeast Pacific near-surface ocean waters may have been as warm approximately 15 000 years B.P. as they now are (Moore, 1973); (c) sea level was rising during the Fraser Glaciation (Clague et al., 1982) and collapse of a marine ice sheet may be very rapid during a rise in sea level even in polar ocean water (Thomas and Bentley, 1978), and (d) late Wisconsin ice appears to have thinned abruptly from the mainland coast towards Queen Charlotte Islands and western Vancouver Island (Clague, 1973; Howes, 1981; and Clague et al., 1982).

After ice had retreated from the coast emergence was rapid and by 10 000 years B.P. sea levels in the Sound were close to those at present (Clague et al., 1982). Sea level continued to drop relative to land, albeit at a slower pace, but may have been up to several metres below present mean sea level during much of middle and late postglacial time (Fladmark, 1974; Andrews and Retherford, 1978; Clague et al., 1982).

Oceanography

Currents, water structure and tidal movements within the Sound have been studied principally by the Pacific Oceanographic Group (1953a,b), Barber (1957a,b, 1958), Canadian Hydrographic Service (1957), Bell (1963) and more recently by Dodimead (1980) and R. Herlinveaux, S. Huggett and R.E. Thomson (in preparation). These studies have (a) defined general tidal flow patterns in the Sound (Canadian Hydrographic Service, 1957), (b) suggested that basin geometry influences movement of local water masses (Bell, 1963; R.E. Thomas, personal communication); (c) indicated that surface water movement in the Sound can be influenced by daily and seasonal wind patterns and by northward coastal water residual flows (R. Herlinveaux, personal communication, 1982); (d) suggested that upwelling occurs in the vicinity of the Scott Islands chain (bounded by Triangle and Cox islands) (R. Herlinveaux, personal communication, 1982); (e) demonstrated that currents 5 m above the bottom can exceed 30 cm/s at depths as great as 325 m (Fig. 5) (R.E. Thomas, personal communication); and (f) detected the movement of deeper halocline water from the open sea into Queen Charlotte Sound. This water intrudes particularly deeply into the Sound (up to Queen Charlotte Strait) during the summer when land drainage is at its maximum and winds are mainly from the northwest, but

there appears to be sufficient year round surface outflow to maintain deep water inflow during all seasons, at least over the deeper shelf areas (Barber, 1958).

Carter (1973) noted that "during the maximum annual storm [along the Canadian west coast] bottom surge velocities in water shallower than 150 m equal or exceed 30 cm/sec" and that this storm surge coupled with local tidal currents may be capable of generating net transport of at least fine sands on the southern Vancouver Island shelf. More recently, Yorath et al. (1979) have shown that storm surges are capable of transporting coarse sands and carbonate shell hash to depths of 105 m on the northwestern Vancouver Island shelf.

Sediment Sources

A review of earlier studies of the character of sedimentation off British Columbia leaves no doubt that glaciers contributed large amounts of sediments to the continental margin during the Pleistocene (Mathews, 1958; Nayudu and Enbysk, 1964; Carter, 1973). Supply of sediment to Queen Charlotte Sound during Holocene time has been sharply reduced because no rivers of consequence flow directly onto the shelf (Tully, 1952) and little sediment is introduced either through Queen Charlotte Strait (Cockbain, 1963) or from the north as the major rivers north of the Sound drain mainly across Dixon Entrance (Crean, 1967; Thomson, 1981). Many sediment-charged rivers on the mainland empty into deep fiords which act as settling basins for much of the land-derived detritus. As material in suspension during periods of highest discharge in typical British Columbia fiords is finer than 49 microns (Pickard and Giovando, 1960), silts are the coarsest sediment that presently can reach the shelf through the fiords. Silt and clay which does reach the shelf must be derived mainly from the large plume of less saline water discharged during the summer from the fiords in the vicinity of the Sea Otter Shoals (Dodimead, 1980).

FIELD AND LABORATORY PROCEDURES

Field

All grab sampling, coring and underwater photography was performed from the following Canadian Forces Auxiliary Vessels at the indicated times: "Laymore" (June 12-17, 1967), "Endeavour" (July 7-13, 1967; August 19-30, 1968 and June 16-21, 1969). Continuous seismic reflection profiles were obtained in 1973 with 16 cm³ and 82 cm³ air gun sources by D.L. Tiffin from the "Endeavour". High resolution profiling was performed during May 30th-June 10th, 1981 from the Canadian Survey Ship "Hudson" with a Huntec (70) Ltd. Deep Tow System (50-500 joules boomer). Side viewing sonar records were acquired on the same cruise with a system developed by Jollymore (1974) and operated by A. Boyce of the Atlantic Geoscience Centre.

Navigation for the 1967-73 cruises was accomplished, depending on weather conditions and distance from shore, by employing various combinations of Decca radar, radio beacon, Loran-A, dead reckoning and detailed bathymetric maps (position accuracy: better than one half sampling interval distance). Navigation for the "Hudson" cruise was accomplished with the integrated BIONAV system employing mainly LORAN C and satellite position fixing (position accuracy: ± 200 m).

Grab samples were collected primarily with a Petersen device although, on occasion, a La Fond-Dietz clam snapper was used. If the sample was a mud, muddy sand or well sorted sand, no more than a litre was retained. Sufficient gravelly sediment was collected to fill at least a 4 litre

container. If insufficient sample was obtained during an earlier cruise attempts were made during subsequent cruises to obtain larger samples. An attempt was made to extract live organisms, especially holothurians, from muddy sediments to prevent them from affecting organic carbon determinations. Two- or four-metre-long barrels having inner diameters of 7.5 cm were used for gravity coring. The floor of the Sound was photographed at eleven locations (of which nine are discussed in this report) with an Edgerton, Germeshausen and Grier camera and strobe unit.

The site at which samples and underwater photographs were obtained for this study are shown in Figure 6. Seismic profile and sonogram trackline locations are displayed in Figure 22.

Laboratory

After having been treated for the removal of organic matter (with H₂O₂) and salt, subsamples were analyzed for grain size (0.5 ϕ sieve interval for all but the mud fraction) according to conventional sieve and pipette techniques (Folk, 1965; Ingram, 1971; Galehouse, 1971). Mean size, standard deviation and skewness were calculated using a computer by the method of moments (McBride, 1971). The computer program defined mid-size-interval points from a linearly interpolated and extrapolated log-probability cumulative frequency curve.

Mineralogy of the <2 μ m fraction of 38 surficial samples and several core subsamples was determined by standard X-ray diffraction techniques (Warshaw et al., 1960; Brindley, 1961a; Kodama and Oinuma, 1963 and Biscaye, 1964). Component clay and other minerals were identified from diagnostic diffraction peaks and responses to various chemical treatments (Weaver, 1958; Bradley and Grim, 1961; Brindley, 1961b; MacEwan, 1961; Walker, 1961; Kodama and Oinuma, 1963 and Biscaye, 1964).

The colour of moist sediments was estimated from the Geological Society of America Rock Colour Chart (1963).

A determination was made of the heavy mineral concentration (expressed as weight per cent) within the 0.177-0.063 mm fraction of best sorted samples with similar mean sizes (ranging from 0.125 to 0.250 mm) using standard bromoform methods (Folk, 1965). Total concentration in a sample was estimated by multiplying the proportion of heavy minerals in the examined size fraction by the proportion of the total sample represented by the analyzed size fraction.

The concentration of coarse, test-free glauconite pellets (the term glauconite is here used as it is defined by Burst (1958)) was determined by point counting a total of 300 grains in the 0.354-0.500 mm size fraction of all samples in which grains of this size were present. Rock fragments, mineral grains, shell material, faecal matter and glauconite pellets were identified. Glauconite pellets were distinguished from faecal pellets by their irregular shape and by their colour (the ovoid faecal pellets tend to be light olive grey (5y 5/2) - pale olive (10y 6/2) while the glauconite is moderate olive brown (5y 4/4)). The weight per cent proportion of 0.354-0.500 mm glauconite in a sample was approximated by multiplying the grain count percentage by the proportion of the entire sample represented by the size fraction analyzed.

The concentration of skeletal carbonate within the sand plus mud fraction of samples was determined with the Chittick Apparatus (Dreimanis, 1962) after being ground finer than 0.5 mm in a mortar and pestle. Noncarbonate organic carbon content concentration was calculated by first determining the total organic carbon content within the sand plus mud fraction of the sample using a Leco Carbon Analyzer and subtracting from this the calculated concentration of skeletal carbon in the sample.




Figure 6. Location of grab and core samples and camera stations.

Foraminifera were extracted from one core and examined by B.E.B. Cameron and C. Rodrigues. The internal structure of cores was examined by means of industrial X-rays. (Film: Kodak type M film with Snofront Screen; Source Power: 4.5 mA and 80KV; Exposure time: two minutes).

RESULTS

Grain-Size Distribution

Sediments within Queen Charlotte Sound are classified into arbitrary mean size - standard deviation - skewness categories and mapped (Fig. 7,8). Figure 8 can be considered only a very generalized representation of the sediment distribution within the Sound, determined entirely from the character of the sediment samples acquired. Bottom photographs and sonograms, which are described in a following section, reveal that the various sediment populations are interrelated in a far more complex fashion.

Sediments from areas of the seafloor shallower than 100 m are virtually free of mud (Fig. 9). A gradual increase in mud content is apparent for most sediment lying between 100 and 140 m. Within this depth range only those samples recovered from the vicinity of the Sea Otter Shoals area,

near the principal present-day source of sediments, have a substantial proportion of mud. Between 140-160 m (which includes the depth range of the inlet mouths) the mud-to-sand ratio increases sharply. Highest mud concentrations were found in sediments lying between 160-200 m, but in approximately three quarters of all samples recovered from depths greater than 160 m (which include those of the outer and northern shelf) mud constitutes less than one third of the sediment.

Bottom Photographs

Ripples evident in photographs taken of the seafloor in different areas of the Sound (Fig. 6) have been classified according to Harm's (1969) scheme. Compass vane and compass in photographs measure 25 cm and 8 cm, respectively.

Sediments at Station 437 (Fig. 10a,b) on Cook Bank near the coast of Vancouver Island are carbonate-rich sands and gravels (Fig. 10a). In Figure 10b both the compass vane and the kelp at the bottom left hand corner of the photo indicate that a current is flowing in a north-northeasterly direction. The particles suspended over the ribbon of kelp at the right of the photo as well as the high energy linguoid ripples suggest this current is capable of sweeping coarse sand and fine gravel size detritus across the seafloor.




Figure 7. Mean size - skewness - sorting categories used to map sediment distribution displayed in Figure 8. Dominant skewness sign displayed by samples in each category indicated by + or -. No sign is indicated in categories D and G where no strong tendency is displayed.




Figure 9. Plot of the ratio of weight per cent mud to weight per cent sand vs sample depth. Numbered dots represent samples collected in the vicinity of the Sea Otter Shoals.




Figure 8. Sediment distribution in Queen Charlotte Sound determined and extrapolated from the character of sediments collected at sites indicated in Figure 6. Local minimum mean grain size indicated in areas having type F sediments.




Figure 10

a.b. Camera station No. 437; Cook Bank (40 m).

- a. *Holothurian*, pelecypod, and white plumed anemone on poorly sorted shelly gravel surface. Carbonate productivity here is one of the highest off western Canada.
- b. Linguoid current ripples have formed in response to high north-south flowing tidal currents.

- c. Camera station No. 267; Goose Island Bank (37 m). Low energy current ripples (oriented east-west) are superimposed on wave generated symmetrical ripples (oriented north-south). Wave and current action is strong enough to sort both sand and gravel.

- d. Camera station No. 311; Goose Island Bank (50 m). Blanket of very well sorted rounded pebbles.

Canadian Hydrographic Service chart 3744 indicates that surface currents in this area can be as high as 150 cm/s. High energy lenticular current ripples also were present at a deeper site on the bank (Station 429) where the bottom is blanketed with very fine to fine (Group D) sands.

Although water depth at Station 435 (Fig. 11d) is similar to that at Station 429, sediments at the former site probably are coarser (Group C sands) as ripples are poorly defined and the dense network of snail trails suggest these sediments are not readily mobilized (Kulm et al., 1975).

The seafloor at Station 267 (Fig. 10c) near the shallowest part of Goose Island Bank, is blanketed with what appear to be two distinct sediment types: sands and a well sorted basal gravel (Fig. 10c). East-west striking, low energy current ripples are superimposed on larger north-south striking, wave generated, symmetrical ripples. Well sorted rounded pebbles (Fig. 10d) without associated sand were photographed at somewhat greater depth on the same bank at Station 311 (these sediments are very similar to those collected at Station 450) (Fig. 6,7,8).




Figure 11

a, b, c. Camera station No. 430; Goose Island Bank (35-40 m).

- a. Indistinct wave generated symmetrical ripples in sand.
- b. Intermixed sand and gravel.

c. Dense pavement of immobile, heavily encrusted gravels.

- d. Camera station No. 435; Cook Bank (73 m). Weak wave generated symmetrical ripples in what probably is medium to coarse sand. Note dense network of snail trails.

Figures 11a,b,c illustrate the extreme changes in sediment character evident over relatively short distances (approximately 3.5 km) at similar depths on Goose Island Bank. Figures 12a,b reveal that the seafloor at Station 391 on Middle Bank is blanketed with debris ranging from boulders to poorly sorted gravelly (Group G) sands. The sand size sediments there consist of a high proportion (up to 51 per cent by weight) of foraminifera and may thus be as mobile as terrigenous sands of the same mean size at shallower depths (Schafer and Prakash, 1968). The north-northeasterly flowing, bottom current, which inclines the crinoid radii in Figure 12a, probably has contributed to the formation of the well defined, wave dominated combined flow ripples in Figure 12b. Photographs of the outer part of Goose Island Trough

(Stations 424 and 427; Fig. 12c,d), in areas where the seabed is blanketed mainly with Group E silts and fine sands, display an abundant fauna and evidence of bioturbation, but no hydraulic bedforms.

Sonograms

The record representing a section of the shallowest part of Goose Island Bank (Fig. 2,3,13A-A') displays alternating bands of high (dark) and low (light) acoustic reflectivity. The bands mainly reflect differences in sediment character as relief is very muted (relief on these Deep-Tow records is an artifact of the system). As coarser sediments tend to reflect acoustic energy more strongly than do finer sediments (Swift




Figure 12

a, b. Camera station No. 391; Middle Bank (130 m).

a. Wave dominated combined flow ripples with lee slopes facing north in foraminifera-rich gravelly sands.

b. Variety of organisms (asteroid, crinoids, and sponges) inhabiting rock strewn bank surface.

c. Camera station No. 424; Goose Island Trough (183 m). Muddy seafloor inhabited by ophiuroids and echinoids.

d. Camera station No. 427; Goose Island Trough (250 m). Muddy seafloor on which are evident burrow holes and trails but no hydraulic bedforms.

and Freeland, 1978), the dark bands may define zones with higher concentrations of gravel (Groups A,B sediments) and lighter bands those areas blanketed mainly with well sorted fine (Group D) sands. Elongated sand patterns displayed in Figure 13A-A' may represent flow parallel "current lineations" (McKinney et al., 1974; Swift and Freeland, 1978) and (or) flow transverse sand patches (Belderson et al., 1972). The darker bands can be considered "deflation zones" (Swift et al., 1978) i.e. areas in which most sand has been swept clear of the basal gravel (Fig. 10c,10d,11c). When the distortion in scale parallel to with respect to that normal to the ship's track is compensated for, the average orientation of the lineations is about 20°T.

The record across outer Intertrough Area (Fig. 22,14) displays crisscrossing grooves varying in width and length and scattered closed depressions. The former probably are relict iceberg furrows (Belderson et al., 1972; King, 1976); the latter possibly are ice-scour craters (McLaren, 1982). The deepest occurrence of a furrow evident on records acquired across the length of Mitchell's Trough is at 200 m at the bend in the trough.

Clay-Size Sediment Mineralogy

Based on the height of the first order basal peaks in X-ray diffractograms (Fig. 15) it is concluded that chlorite and smectite are the dominant phyllosilicates in bottom sediments. Illite tends to be somewhat less abundant and only trace amounts of kaolinite and vermiculite (or mixed layer clays) are present. Small amounts of quartz, feldspar and amphiboles probably also are present.

Heavy Minerals

Extraction of heavy minerals from best sorted sands having similar mean size revealed that the six samples with the highest concentrations, numbers 27 (91 m), 63 (122 m), 74 (102 m), 180 (82 m), 310 (92 m) and 312 (174 m) (Fig. 16) have mean sizes ranging from 0.145 to 0.205 mm. All but the last of these samples fall within a 40 m depth range centred about 100 m. Three of the samples lie near well defined or muted canyons on the northern margin of Cook Bank. The other three lie along the southwestern margin and slope of Goose Island Bank. The weight per cent proportion of magnetite in the heavy mineral fraction of the above samples ranges from six and eight per cent (numbers 312, 74, 27) to 29 per cent (number 63).

Glaucinite

Glaucinite was found either enclosed in foraminiferal tests or in a free state (in the latter case it often retains the form of the host test) (Fig. 17). Although glauconite is present in most sediment groups on the shelf (Fig. 16), few if any of the sediments have large concentrations of the coarsest, test-free and, presumably, better developed glauconite. However, a high proportion of the shelf edge - upper slope sediments have a relatively high concentration of these glauconite pellets, an association suggested earlier by the work of Murray and MacKintosh (1968). The genesis of what appears to be similar test-associated glauconitic matter on the Norwegian shelf has been described by Bjerkli and Østmo-Saeter (1973).




Figure 13. Sonogram and Deep-Tow seismic records of shallowest part of Goose Island Bank. Maximum thickness of sand lenses in Deep-Tow section cd: ~2 m. Refer to text for interpretation (GSC Line 81-1 to 2).

B'

B



Figure 14. Sonogram of central Intertrough Area displaying relict ice furrows (GSC Line 81-2 to 3).




Figure 15

X-ray diffractograms illustrating representative responses of fractions of bottom samples finer than $2\ \mu\text{m}$ to various analytical treatments (Cu K α radiation).




Figure 16. Glauconite pellet and heavy mineral distribution. Numbered black dots indicate total concentration (weight per cent) of 0.177-0.063 mm; heavy minerals in best sorted sands with similar mean sizes (0.45-0.205 mm). Numbered open circles indicate total concentration (weight per cent) of coarse (0.345-0.250 mm) glauconite pellets in the mud plus sand fraction. Unnumbered open circles indicate sites where coarse glauconite was present but constituted <1% by weight of mud plus sand fraction.




Figure 17

The photo at left displays a variety of semi-enclosed and test-free glauconite particles found in Queen Charlotte Sound sediments. The tests and pellets range in size from 0.354 to 1.00 mm. On the right are two samples of what are identified as test-free glauconite pellets ranging in size from 0.354 to 0.500 mm.

Gravity Cores

Core 448 obtained from that portion of Goose Island Trough northwest of the Sea Otter Shoals (Fig. 6) exhibits a general downward decrease in grain size (but is essentially a clayey silt throughout) and organic content (Fig. 19). Colour grades from greyish olive green (5GY 3/2) at the top (Fig. 20A) to dusky yellow green (5GY 5/2) at the bottom. X-ray photographs revealed distinct laminations only in the upper third of the 275 m long core (not illustrated). The bottom two thirds appears to be a massive deposit with a few small molluscan shells (Fig. 21).

Core 484 from Mitchell's Trough flats (Fig. 6) is capped with an olive grey (5Y 3/2) slightly clayey silt-sand with low to moderate amounts of organic matter, somewhat similar to that in core 448 (Fig. 19). These sediments overlie, at a sharp contact, 30 cm of intercalated dark greenish grey (5G 4/1) clayey silt and material similar to that at the top of the core (Fig. 20, core C₁). The remaining length of the core is entirely dark greenish grey clayey silt (Fig. 20, core C₂). Foraminiferal assemblages dominated by *Epistominella* sp. obtained from the lower sediments suggest they were deposited in a late Quaternary open shelf environment (B.E.B. Cameron, personal communication, 1974). The bottom three quarters of the core is more silty and has less organic matter than the sediments at equivalent depths in core 448 (Fig. 19).

The uppermost sediments in core 492 from the outer part of Mitchell's Trough (Fig. 6) are similar in grain size and colour to those at the top of core 484 (Fig. 19). These overlie at a sharp contact (Fig. 20, core B) dark greenish grey sandy clayey silt similar to, but somewhat coarser than, that in the lower unit of core 484. X-ray images (Fig. 21) indicate the greyish olive green sandy silt at the top of the core is laminated whereas the dark greenish grey finer sediment in the lower portion of the core may be massive.

Core 520 from the outer part of Goose Island Trough (Fig. 6) exhibits a downward increase in grain size from clayey silty sand to clayey sandy silt (Fig. 19). At equivalent depths noncarbonate organic carbon concentrations generally are higher than in cores 492 and 484, but less than in core 448. On average, carbonate concentration is higher in core 520 than in any other core. Colour grades from olive grey (5Y 3/2) to greyish olive (10Y 4/2). X-ray photographs (Fig. 21) reveal that coarse laminations and, possibly, cross-bedding are present to a depth of 60 cm in the core below which the sediments are very finely laminated or massive.

Proportions of chlorite and illite tend to increase slightly with depth at the expense of smectite in olive grey and greyish olive green sediments in cores 520 and 448, but are distinctly dominant in the dark greenish grey sediments below the unconformities in cores 484 and 492.

In summary, sediment structure, texture and composition changes gradually downward in Goose Island Trough cores whereas deeper deposits in Mitchell's Trough cores are separated from surface muds, accumulating under present conditions by a sharp, possibly erosional, unconformity and are distinguishable from them by all the following characteristics: greenish grey vs olive grey colour, finer grain size, lower organic but higher chlorite and illite contents and finer or less well-defined laminations.

Continuous Seismic Profiles

Line drawing interpretations of selected seismic records presented in this section delineate major reflectors, identify the uppermost laterally continuous angular unconformity (heavy line) and, in certain cases, parallel bedded or acoustically transparent sediments capping the stratigraphic section (stippled). Profile locations are displayed in Figure 22. Sediment thicknesses are calculated assuming a sound velocity of 1500 m/s.



Figure 20. Photographs of representative sections of cores obtained in Queen Charlotte Sound troughs.

A. Core No. 448. Inner Goose Island Trough. Uppermost 50 cm of core. The entire core displays fairly uniform colour (greyish olive to dusky yellow green) characteristic of muds lying on present seafloor.

B. Core No. 492. Outer Mitchell's Trough. Section of core displays major unconformity (29-30 m from top) between upper olive grey sand-silt and lower, predominantly dark greenish grey, clayey sandy silt. Whitish patch near contact represents remains of a bivalve.

C₁ and C₂. Core No. 484. Central Mitchell's Trough. C₁ represents uppermost and C₂ lowermost section of 275 cm core. Note similarity between this core and core No. 492 at contact between darker, coarser, upper sediments and lighter, finer, lower sediments. In this core the unconformity lies 17 cm from top of core. The protrusion of darker sediment into underlying silts probably is an infilled burrow hole.

Profile A-A' (Fig. 23). Folded (?) units which lie adjacent to and, possibly, under Middle Bank are unconformably overlain by generally flat lying beds which, in turn, are unconformably overlain by laterally discontinuous sediments (~100 m thick) which appear to cap this section and extend from the trough onto adjacent banks. The upper unconformity delineates at least two distinct V-forms (scour and fill features?) about 30-40 m deep and 1-2 km wide. The Middle Bank wall of the trough appears primarily to reflect an erosional origin whereas the Goose Island Bank wall is more likely an accretionary feature. Another record (not illustrated), however, displays what appear to be truncated units on this wall approximately 15 km to the southwest of this site.




Figure 21. Representative X-ray images of sections of cores obtained in Queen Charlotte Sound. Deeper parts of core 448 (inner Goose Island Trough) are massive. Core 492 (outer Mitchell's Trough) displays well developed coarse laminations and what appears to be crossbedding in the sandy sediments capping the core. Core 520 (outer Goose Island Trough) displays crossbedding at its top which grades into flat lying finer laminations which in turn grade into a very finely laminated or massive deposit at the bottom.




Figure 22. Index map of locations of seismic profile tracklines (heavy lines) and side viewing sonar tracklines (dotted lines). Contours in metres.

Profile B-B' (Fig. 24): The apparently folded "bedrock" core evident in the preceding profile is here draped by a thick succession of parallel beds which extends down Mitchell's Trough and under the slope. These beds are unconformably overlain by irregularly stratified sediments which are as much as 75 m thick (or ~100 m if a sound velocity of 2000 m/s. is assumed). Deep-tow seismic records indicate that horizontally bedded, transparent material capping the sequence in places along the outer trough is no more than about 5 m thick. The shelf edge sill appears to consist of a core of the deeper parallel bedded strata overlain with a thin veneer of what is probably poorly sorted detritus. Similar stratigraphic relationships are displayed in a seismic profile along the axis of Goose Island Trough (not illustrated).

Profile C-C' (Fig. 25): This profile is an extension of Profile B-B'. Deeper parallel bedded units are unconformably overlain by a sequence of laterally discontinuous layers which are capped by apparently flat lying, relatively thin bedded deposits having a maximum thickness of about 20 m.

Profile D-D' (Fig. 26): On the mainland (left) end of the profile what probably is crystalline bedrock outcrops on the seafloor and slopes below the surface of Mitchell's Trough flats. Approximately half way across the trough a wedge of gently sloping, parallel-bedded sediments abuts this surface and, in places, may outcrop on the trough floor. Above an erosional surface extending across both the crystalline and sedimentary sequences lie what appear to be mainly horizontally bedded sediments which now form the floor of the trough. On a profile extending from Goose Island to the mainland (not illustrated) the contact between crystalline and sedimentary rocks also underlies the trough floor.

Profile E-E' (Fig. 27): The domed feature at the base of this section probably is a subbottom extension of Sea Otter Shoals crystalline rocks. A thick succession of seaward

dipping sediments over this bedrock core is in unconformable contact with overlying irregularly bedded sediments which form a continuous cover over this part of the shelf. At the centre of this section the sedimentary column is capped by parallel bedded sediments having a maximum thickness of about 35 m.

Profiles F-F' and G-G' (Fig. 28): The air-gun record (F-F') normal to the axis of Moresby Trough indicates that parallel bedded deposits are concentrated in the central depression which is bounded on the east by a relatively steep scarp of either erosive or tectonic origin. The deep-tow seismic record depicting the central area along a trackline (G-G') oriented obliquely to the thalweg of the trough reveals (a) what appear to be slump blocks at the base of the scarp, (b) truncated sediment layers at the floor of the deepest trench within the trough and (c) an upper sequence of sediments (maximum thickness ~30 m) to the west of the trench with bedding which is laterally continuous and uniformly thick over several kilometres. These sediments overlie, in places unconformably, a similar sequence of strata which may extend beneath the lower, essentially featureless, terrace on the northwestern flank of the trough. These laterally continuous deposits have been derived from the west-northwest and have ponded behind and overridden a now buried rise.

Profile H-H' (Fig. 29): This record suggests bank top sediments are prograding across the northernmost flank of Goose Island Bank to a depth of 170 m onto the irregularly bedded deposits which form the floor of the Intertrough Area. An acoustically discontinuous irregular surface underlies the disjunctly bedded sediments. Other records indicate a comparable wedge of sediments is also accumulating off the southeastern margin of the bank but extends only to 100 m depth.




Figure 23. Air-gun seismic profile A-A' across central Mitchell's Trough. Refer to text for interpretation (GSC Line 73-13).




Figure 24. Air-gun seismic profile B-B' along axis of outer Mitchell's Trough. Refer to text for interpretation (GSC Line 73-12).

Profile I-I' (Fig. 30): Sediments probably derived from Cook Bank are shown here to have ponded behind and overridden a now buried rise or ridge. Much of the sequence of sediments may represent glacial outwash deposits. The relatively thin, chaotic bedding of some of the sediments on the flank of Cook Bank, however, suggests slope failure and concomitant turbidity flows also may have contributed to sediment deposition in Goose Island Trough.

Profile J-J' (Fig. 31): Irregularly bedded sediments having a maximum thickness of about 50 m unconformably overlie older deposits. The sediment column is capped by parallel bedded sediments which have accumulated asymmetrically across the trough floor suggesting they have been derived primarily from the north.

In summary, the seismic records appear to indicate that:

1. The present shelf and slope surface has developed on a sequence of sediments which prograded westward over what probably is crystalline bedrock on the inner shelf and sedimentary bedrock on the outer shelf (Fig. 24).
2. During a major glaciation the shelf was scoured across its entire width (Fig. 24) and to depths as great as 75-100 m below the present floor of the part of troughs at the centre of the shelf (Fig. 23).
3. Sills at the shelf edge are erosional remnants of possibly pre-Pleistocene or early Pleistocene deposits (Fig. 24). Mid-shelf ridges are, in places, continuous with irregularly bedded (drift?) deposits (Fig. 25) overlying parallel bedded sediments.
4. Sections of troughs parallel to the mainland coast lie along the contact between crystalline coastal rocks and thick sediments underlying the shelf (Fig. 26).




Figure 25. Air-gun seismic profile C-C' at the bend in Mitchell's Trough. Refer to text for interpretation (GSC Line 73-12).




Figure 26. Air-gun seismic profile D-D' across central Mitchell's Trough flats. Refer to text for interpretation (GSC Line 73-4).




Figure 27

Air-gun seismic profile E-E' along axis of Goose Island Trough north of Sea Otter Shoals. Refer to text for interpretation (GSC Line 73-37).

5. The lower terrace on the northwest flank of Moresby Trough is underlain by a sequence of uniformly bedded outwash deposits (Fig. 28).
6. Eroded sedimentary strata lie at or near the sediment-water interface in the deepest basin within Moresby Trough (Fig. 28).
7. Troughs (with the possible exception of Moresby Trough) are not fault bounded nor do they lie along identifiable fault lineations (Fig. 23, 26, 29, 30, 31).
8. Now-infilled shallow troughs linked Milbanke Sound across the Intertrough Area to Mitchell's Trough (Fig. 29).
9. Outer trough walls of Goose Island Trough are more consistently accretionary in character than those of the more northerly troughs (Fig. 23, 28, 30).
10. Thickest accumulations of localized parallel bedded (mud?) deposits (~30 m thick) underlie the lower terrace on the northwestern flank of Moresby Trough and form the uppermost deposits at the bend in Mitchell's Trough and in Goose Island Trough north of the Sea Otter Shoals (Fig. 25, 27, 28). If it is assumed that these deposits have been accumulating from 13-15,000 years B.P. to the present, the average sedimentation rate at these isolated sites is 2-3 mm/a. (Even these few sites however, may not all be major present day sediment sinks.)
11. Currents are sweeping sediments to the north and south of the eastern side (Fig. 29) and, possibly, to the southwest of the western part of Goose Island Bank (Fig. 31).




Figure 28. Air-gun seismic profile F-F' across outer Moresby Trough. Refer to text for interpretation (GSC Line 73-5). Deep-Tow high resolution seismic profile G-G' obliquely crosses axis of outer Moresby Trough. Refer to text for interpretation. (GSC Line 81-22 to 21).



Figure 29. Deep-Tow high resolution seismic profile H-H' across the northeastern margin of Goose Island Bank. Refer to text for interpretation (GSC Line 81-1 to 2).




Figure 30. Deep-Tow high resolution seismic profile I-I' across the southern flank of outer Goose Island Trough. Refer to text for interpretation. (GSC Line 81-4 to 5).




Figure 31. Deep-Tow seismic profile J-J' across the northern flank of outer Goose Island Trough. Refer to text for interpretation (GSC Line 81-5 to 6).

DISCUSSION

Structural/Stratigraphic Influence on Shelf Physiography

The gap between the Queen Charlotte Islands and Vancouver Island offered the only unobstructed passage for Pleistocene ice spilling off the central Coast Mountains. This breach in the Insular Belt must have determined to a large extent the general northeast-southwest trend of the troughs in Queen Charlotte Sound. The specific orientation and location of different segments of the troughs, however, probably were defined by local structural and stratigraphic lineations such as those, for example, that are considered to have influenced the orientation of "marginal channels" off glaciated coasts (Holstedahl, 1958, 1970; Grant, 1972).

These geomorphic features lie at high angles to the dominant direction of ice movement (in the Sound it is assumed to have been from the northeast to the southwest). Their orientation is considered to reflect selective glacial erosion of a fault zone and (or) the contact between crystalline coastal rocks and the thick sedimentary deposits on the shelf (Holstedahl, 1970; Grant, 1972). Examples in Queen Charlotte Sound are the section of Goose Island Trough between Goose Island Bank and the mainland and the flats portion of Mitchell's Trough which are oriented normal to the presumed principal directions of local ice flow and lie along stratigraphic contacts. There is no evidence to indicate that faulting has influenced the glacial excavation of these trough sections.

Factors which may have been instrumental in defining the specific location and orientation of the troughs which generally parallel the direction of ice flow are somewhat more obscure. Cook Trough parallels lineaments on both the adjacent mainland and northern Vancouver Island and may lie along a deep fault or stratigraphic contact. Outer Moresby Trough and outer Goose Island Trough may lie along the contact between more indurated Tertiary and less indurated younger deposits on the shelf. The low level of seismicity of the Queen Charlotte Sound area would suggest that even if future surveys confirm that Moresby Trough does lie along a fault it probably is not the locus of major present motion.

As crystalline rock probably underlies the Intertrough Area, it is possible that this broad area of low relief is a submerged and now buried extension of the Milbanke Strandflat. Submerged strandflats, in fact, have been recognized off other glaciated coasts (Holstedahl, 1958, 1970).

Ice Contact and Proglacial Features

The ridges crossing troughs are probably recessional or end moraines deposited at the terminus of a grounded ice lobe or below an ice shelf a relatively short distance beyond the grounding line (Barnett and Holdsworth, 1974; Fillon, 1975). King et al. (1972) suggested that sills lying along a shelf edge develop as a consequence of fracturing and calving of glaciers where water depth increases abruptly, presumably because of the diminished erosive capacity of ice and (or) by the release of enclosed detritus under these circumstances.

The complex of features at the intersection of Cook Trough and Goose Island Trough is similar to the morainal and ice contact landforms (kames, kettles, esker?) associated with stagnant or receding glaciers on land (Flint, 1957). Consequently, the anomalously deep, well sorted, fine sands in Goose Island Trough immediately to the west of this area possibly represent relict outwash fan deposits. As the ridges at the bend in Mitchell's Trough are the same depth as the

features at the intersection of Cook and Goose Island troughs, they may represent contemporaneous stagnation sites of receding grounded glaciers.

The morphology and sediment character of the Intertrough Area, a part of the Sound which still displays evidence of intense scouring by icebergs, resembles that found on areas of low relief, both on land and in the sea, which have been glaciated relatively recently (Holmes, 1965; refer to Figures 500 and 504; Flint, 1957; Fillon, 1976; Van der Meer et al., 1976). Ice receding across the Intertrough Area flushed fine sediment into the adjacent trough to form Mitchell's Trough flats (the area less than 220 m deep). Middle Bank inhibited dispersal of this glacial outwash to Moresby Trough. What detritus did accumulate in this sediment-starved basin seems to have been derived primarily from ice which approached and retreated across western Hecate Strait. This ice probably stagnated on or over the shallower hummocky terrace on the northwestern flank of Moresby Trough and formed the featureless deeper terrace which is underlain by uniformly bedded outwash deposits.

Outwash from ice which had stagnated at the muted sill which extends between Goose Island Bank and the Intertrough Area may have buried a shallow trough which linked Milbanke Sound to Mitchell's Trough and led to the formation of (a) the low, long, spit-like ridge (at 120 m) extending across what may be a fault controlled embayment on the north side of Goose Island Bank, (b) the shallow elongate canyon just north of the ridge and (c) the parallel bedded deposits nestled in depressions in drift at the head of the transverse section of Mitchell's Trough.

Formation of Troughs and Banks

The formation of troughs on the West Antarctic shelf (e.g. Hughes, 1973; Thomas and Bentley, 1978) and Labrador shelf (Fillon, 1975) has been attributed at least in part to scouring by ice streams within submerged grounded ice sheets. The same mechanism can also explain the formation of troughs in Queen Charlotte Sound where the relief is, and probably was at the time of excavation, too muted to have contained isolated submerged valley glaciers which extended across the shelf. Formation of the troughs must postdate that of the aprons of sediment at the southwest of Goose Island and Middle Banks as troughs dissect the aprons. Consequently, the aprons may be outwash deposits laid down ahead of an advancing ice sheet and reworked by waves and currents.

When ice streams retreated along troughs they mantled the previously scoured surface with glacial debris including the identified ice contact and proglacial deposits. As ice sheets are expected to retreat more slowly on higher ground where they may remain grounded longer than in adjacent troughs (Thomas and Bentley, 1978) masses of ice may have been stranded on bank tops after trough ice had receded past the banks. Meltwater streams coursing through caverns at the base of this ice probably formed the channels on Cook and Goose Island banks.

Glacial sediments constituting the banks accumulated on bedrock platforms (e.g. the northward extension of Nahwitti Lowland which probably underlies Cook Bank and at the top of the parallel bedded units which underlie Middle Bank). The height of such platforms and the volume of glacial debris deposited at any one site largely determined the highly discordant depths of the banks. The gentle seaward concavity of the terraced surface of Middle and Goose Island banks suggests it also reflects the effects of surf and (or) wave scour.

Extent of Ice Grounding During the Fraser Glaciation

Although no deposits on the shelf have been dated, certain lines of evidence suggest Fraser ice not only could have been grounded on the shelf but that grounding may have extended across the entire shelf:

1. If it is accepted that during the Fraser Glaciation (a) the outer shelf underwent little or no depression (and may as a result have been shallower or no deeper than the isostatically depressed inner shelf) and (b) sea level was lowered ~75–100 m, an ice thickness of probably no more than 300–400 m (1.1 times water depth) was required for an ice sheet to be grounded at the depth of the base of the drift deposits in Mitchell's Trough (75–100 m below the seafloor of the central shelf but less than that at the outer shelf). This ice thickness is less than half of the minimum probably required to induce the estimated 120+ m depression along the northwestern coast of Vancouver Island if the ratio of glacioisostatic depression to ice thickness of about 0.10 to 0.15 presumed to have prevailed on the mainland is applied to this area.
2. Icebergs from what is assumed to be the last major glacial event scoured the seafloor to depths at least as great as 200 m on the present inner half of the shelf.
3. Margins of the outer part of the troughs are still well defined, shelf edge sills are still apparent and Moresby Trough has a sharply irregular relief. If such features are pre-Fraser in age they would have had to survive blanketing by debris discharged during the retreat of a penultimate glaciation and during the full cycle of Fraser Glaciation. Furthermore these features were exposed to tidal current and wave erosion and sediment redistribution during successive regressions and transgressions of the sea that have occurred during the >67,000 years (Clague, personal communication, 1982) that elapsed since the penultimate glaciation.

An alternate possibility, that grounded ice of the Fraser Glaciation extended only part-way across the shelf within troughs, is supported by these considerations:

1. Sea level was rising during the last major glaciation and local water temperatures were nonpolar and may well have been similar to those at present.
2. Well-preserved deposits considered to have an ice front or ice contact origin (including outwash) are concentrated on the central and inner shelf.
3. Relief is more muted and depths generally are shallower toward the mainland and Vancouver Island coasts. This suggests that deposition from the last phase of glaciation was limited largely to that part of the shelf closest to major icefields.

A third possibility is that grounded ice reached the shelf edge but retreated rapidly to mid-shelf from whence it more gradually made its final withdrawal.

Apparently abrupt thinning of the ice front in the vicinity of Queen Charlotte Sound and the relatively short duration of the Fraser Glaciation favour the second alternative. Persistence of the glaciogenic physiographic features on the outer shelf if they were formed by pre-Fraser ice may have been promoted by gravel armouring of the seafloor and very limited sediment discharge by Fraser ice.

Impact of Lower Sea Levels

Intense tidal currents which swept over Cook Bank after ice had retreated or was no longer grounded but before the sea attained its present level, may have scoured the very gently inclined valleys at the margins of the bank. It is not

readily apparent why these are not equally well developed valleys on the other banks, especially, the shallower Goose Island Bank. Cook Bank valleys may have been at least partly scoured in bedrock and, consequently, may be relatively stable features.

Turbulent tidal flows and wave action (Evans, 1939; Rao, 1957; Dubois, 1972; May, 1973) on Cook and Goose Island banks when sea level was lower initiated the sorting of sediments and contributed to the production of the deposits rich in heavy minerals now found at or near 100 m at the margins of these banks.

Curry (1969) suggested that the depth of the break in slope of continental shelves has been mainly controlled (barring post-Pleistocene isostatic movements) by shallow water processes active when sea level was 100–150 m lower because most shelf breaks lie within this depth range. As the constructional parts of the Queen Charlotte Sound shelf break lie at twice this depth and there is no reason to expect that the shelf edge has undergone depression since the Late Wisconsin, mechanisms other than shallow water oceanographic processes must have shaped its character. If we consider (a) that bottom photographs reveal vigorous sediment transport to depths of 130 m, (b) that mud is only a minor component of the majority of sediments on the outer shelf even at depths greater than 160 m, (c) that near bottom currents at 255 m can exceed 30 cm/s and (d) that cores obtained at or near 300 m near the shelf edge reveal what appear to be crossbedded, sandy, near-surface sediments, it is obvious that oceanographic processes are now, or were in a very recent past, capable of mobilizing at least fine sands on the seafloor over most of the outer shelf.

The evidence from this shelf coupled with what is known of shelf edge dynamics in other areas (Southard and Stanley, 1976) make it reasonable to propose, then, that when Late Wisconsin ice (which probably did not markedly depress the outer shelf) was flushing sediment across the Sound and sea level was as much as 100 m lower, bottom turbulence probably was high enough over the site of the present shelf edge to mobilize at least sands and muds and permit continuous accumulation of material mainly below that depth.

Present Sediment Dispersal and Deposition

The bulk of new terrigenous detritus supplied to Queen Charlotte Sound is transported during the summer in surface and near-surface waters from the mouths of fiords in the vicinity of Sea Otter Shoals. The Coriolis effect coupled with a northward coastal water residual flow tends to direct this silt-charged water to the right as it enters the Sound, whereas prevailing northwesterly winds move near-surface waters to the southwest. Examination of surficial sediment distribution, cores and seismic records suggests that the greatest lateral and vertical accumulation of muds having a character similar to that of muds on the present seabed lie in inner Goose Island Trough north of Sea Otter shoals and not in Cook Trough suggesting that the Coriolis effect and the coastal flow more strongly influence the dispersal direction of suspended sediment newly discharged into the Sound than do northwesterly winds. Some mud, which gradually settles through the water column as the meltwater plume crosses the shelf, is probably carried back to the inner shelf floor by advective transport (McCave, 1972) in the colder, more saline bottom waters which most deeply intrude the Sound during the summer.

The location of the major site of present sediment discharge into the Sound, water circulation patterns, and the intensity of wave and current activity on the shelf have insured that (a) mud is not accumulating at depths shallower than 140 m (except near the southeastern coast in the path of

the plume of less saline water) and (b) for any given depth surficial sediments generally are finer on the more sheltered inner and southern parts of troughs and or inlet mouths. Although high sediment transport (Hein et al., 1974) may have helped maintain test-free glauconite at low concentrations on the shelf, that it is present at all probably is a reflection of the generally slow sedimentation rates (Degens, 1965). The somewhat higher concentrations found in muds on the slope indicate that even fine sediment there is being deposited very slowly (Degens, 1965; Hein et al., 1974).

Surficial muds in the Sound have a higher proportion of smectite than the subbottom greenish grey clay size sediments or the muds in the Taku Estuary (Kunze et al., 1968) and Glacier Bay (O'Brien and Burrell, 1970) in southeastern Alaska which are derived from present day glacial abrasion of rocks similar to those on the mainland adjacent to Queen Charlotte Sound. This suggests that the more temperate climate conditions which now prevail immediately south of Alaska encourage chemical "stripping" of chlorite, hornblende and mica, leading to the formation of smectite (Weaver, 1958; Keller, 1970).

The gross inverse relationship between noncarbonate organic carbon concentration and grain size of enclosing sediments could be governed by a number of conditions: (a) coarser sediments are more permeable and thus permit organic matter to be oxidized rapidly, (b) particulate organic matter tends to deposit together with fine grained clastic sediments and (c) clay size particles enriched in clay minerals have large capacities to adsorb organic molecules (Bader, 1962; Thomas, 1969 and Naidu et al., 1975).

The tendency for noncarbonate organic carbon to decrease gradually with depth in a core may indicate a tendency for organic matter to oxidize after burial. The abrupt change in organic content at the sharp contact between the surface sediments and the dark greenish grey mud more likely reflects a fundamental change in the nature of the sediments and mode of sedimentation. The virtual lack of particulate organic matter in the underlying sediments may be due to the low clay mineral content of glacial flour or it may indicate that these muds were deposited too rapidly to allow enough time for substantial organic adsorption and development of a rich bottom fauna. It is possible these muds were deposited under ice but the absence of drop stones and presence of *Epistominella* would argue against this.

The highest concentration of well sorted sediments falls within the 2-3 ϕ size range. Inman (1963) has established that sands of this size have the lowest threshold friction velocities of any sediment. Hamilton et al. (1980) have proposed that "repeated selective entrainment by tidal currents in a virtually closed sediment system [as exists on Queen Charlotte Sound banks]... lead to the formation of such well sorted mobile sands (it is possible, however, that considerable sorting may already have been achieved, if, as probably is the case, these sediments were deposited as outwash). The above authors have suggested that the fine sand population will gradually blanket the seafloor and inhibit mobilization of coarser grades which ambient currents would otherwise be capable of transporting. It can be expected, however, that fine sands will eventually be swept clear of the banks by tidal flows which are known to induce unidirectional sediment transport (Off, 1963; Houbolt, 1968; Belderson and Kenyon, 1969; Kenyon and Stride, 1970; Ludwick, 1974; and Luternauer, 1976, 1978).

The distribution of best sorted fine to very fine sand and the nature and orientation of ripples suggest tidal currents are moving this sediment from Cook Bank onto its northern flank. Photographs of the sands on the eastern edge of this bank reveal a dense network of snail trails. These features probably are preserved because these sediments

generally are coarser and are not likely to be as frequently mobilized as finer sands on which no evidence of trails is found. The less frequent or intense abrasion which these sediments probably undergo may also explain why quartz grains within these sands retain a yellow brown, presumably iron oxide, coating. The coating may have been acquired in a subaerial beach environment (Dolan, 1970) or in a river (Judd et al., 1970) or even in their present environment (Swift and Boehmer, 1972). Swift (1969) and Emery and Niino (1963) have stressed the importance of subaerial exposure in the formation of iron-stained shelf sands, but the critical factor governing the formation of such a stain is the availability of ample amounts of magnetite (which is abundant in these sands) and (or) ilmenite (Miller and Folk, 1955) which can alter readily to limonite under oxidizing conditions (Ollier, 1969).

The distribution of best sorted sands on Goose Island Bank, the character and orientation of bedforms evident in photographs, the orientation of what may be flow parallel "current lineations" and the cross sectional geometry of deposits at the margins of the bank revealed in seismic profiles, strongly indicate that fine sands are being washed by the combined action of waves and tidal currents from the sandy gravels and gravelly sands (Stanley, 1968) mainly towards the north and southwest. The morphologic character of Goose Island Bank is also compatible with such a dispersal pattern: slopes are very gentle to the north and southwest but relatively abrupt toward the south and southeast. Yellowed sands and snail tracks are also evident on this bank and probably occur where sediments are transported only under storm conditions or by spring tidal currents. The glauconite on the bank probably formed in foraminiferal tests which were subsequently split open and carried off in the fashion described by Keller and Richards (1967).

The area on Middle Bank covered by well sorted fine sand appears very small when compared to areas mantled by the same sediment type on Goose Island and Cook banks. This probably reflects the lower ambient wave energies on this, the deepest, bank. As is the case for the other banks, sands appear to be washed to the north and south.

The broadest terraces at the tops of banks all lie above 140 m. Goose Island Bank is the shallowest bank and probably is undergoing most active planation. Cook Bank probably is also being eroded, but to a lesser extent because it is somewhat sheltered from Pacific storm waves and swells by the Scott Islands. The presence of wave dominated combined flow ripples on Middle Bank suggest it is a zone of intermittent erosion occurring mainly during storms. On both Cook Bank and Goose Island Bank foraminiferal tests are very sparse. On much of Middle Bank tests are extremely abundant, although they tend to be concentrated in very poorly sorted sandy gravels. Schafer and Prakash (1968) have found that "the size of Foraminiferal tests and the possible entrapment of air or some other gas makes them most susceptible to erosion and transport by currents. This is especially true of abandoned foraminiferal tests which are no longer inhabited by the living animal." Currents are probably strong enough to sweep forams off Cook and Goose Island banks. On Middle Bank they may settle out when weather conditions are less severe. During a storm foraminiferal tests are washed from better sorted sands on Middle Bank but are not as readily dislodged from poorly sorted sediments trapped between the boulders on the bank surface.

The recorded concentrations of skeletal carbonate in Queen Charlotte Sound lend further support to earlier studies that indicate that abundant carbonate-rich sediments may develop at high latitudes where the supply of terrigenous detritus is low (Chave, 1967; Hoskin and Nelson, 1969; Milliman, 1971; Muller and Milliman, 1973). On Cook Bank upwelling of nutrient-rich slope waters probably has further contributed to the accumulation of coarse shell hash deposits.

Environmental and Economic Considerations

1. Information on the character of bottom materials presented in this report coupled with available data on the occurrence of specific ground fish (Westheim, 1974) may identify possible substrate/fish associations which warrant further investigations. Examination of bottom photographs and sediment distribution can contribute to more precise mapping of ground which can be trawled without equipment damage and may help identify ways in which trawling equipment may be redesigned to render it more suitable for operation in local waters.
2. Information included herein about the nature of surficial sediments, the extent and water depth to which sediments are mobilized, and the location and size of possible faults in the Sound will have to be considered if and when drilling platforms are being designed for hydrocarbon exploration in the Sound.
3. Pollutants released in the course of hydrocarbon exploration or exploitation which settle to the seafloor probably would collect in the inner parts of troughs and inlet mouths, be adsorbed in the smectite rich sediments and could, potentially, disrupt benthic deposit feeders (Hatcher and Segar, 1976; Hein et al., 1979). Drilling and production waste falling onto sand-pebbly bank surfaces probably would not seriously impair this habitat because of the mobility (Anderson et al., 1981) and abrasiveness of these materials. Hydrocarbons trapped within boulder beds, however, may persist substantially longer and affect sessile organisms and ground fish.
4. The best sorted, nonencrusted finer gravels and clean sands best suited for construction materials are most concentrated on the eastern, shallower part of Goose Island Bank. The most extensive skeletal carbonate beds, a potential source of lime for use in the manufacture of cement, extend for several tens of kilometres in a 10-15 km wide band adjacent to northern Vancouver Island and the Scott Island chain. The highest concentration of heavy mineral rich "black sands" are found at or near the 100 m depth on northern Cook Bank. What are probably equivalent deposits at the mouth of the Nahwitti River, which flows on to Cook Bank, have been described by the British Columbia Department of Mines and Petroleum Resources (Anonymous, 1980). It is noted that these deposits contain gold which, however (a) is very fine, (b) is only sufficiently concentrated to be worked after certain storms and (c) is associated with high concentrations of magnetite.

CONCLUSIONS

1. The continental shelf and slope surface of Queen Charlotte Sound has formed on sediments which prograde mainly from the mainland coast over bedrock cores.
2. Some inshore sections of troughs which can be considered "marginal channels" lie at or near the contact between crystalline coastal rocks and offshore sedimentary deposits.
3. Moresby, the deepest trough is the only feature which may lie along a fault active during the late Quaternary.
4. Partially submerged grounded ice sheets enclosing ice streams sculpted morphologic features on the shelf and strongly influenced the present sedimentary framework.
5. Partially submerged ice having a thickness of the same order of magnitude as that suspected to have blanketed the western part of northern Vancouver Island during the late Wisconsin Fraser Glaciation probably was capable of grounding on the outer shelf to the depth of the uppermost laterally continuous unconformity (which is 75-100 m below the trough floor of the centre of the shelf) if it is assumed the submerged shelf was not markedly depressed.
6. Much of the material constituting the slopes of Goose Island and Middle Bank between the bank tops and continental slope probably was deposited ahead of the ice sheet which scoured the troughs. These sediments were subsequently reworked by late Pleistocene seas.
7. The following ice contact and proglacial features or deposits are present on the shelf: terminal and/or recessional moraines, kame and kettle topography, an esker, and outwash sands and muds.
8. The progressively greater depth and increased morphologic irregularity of the outer parts of more northerly troughs suggests that deposition by late Wisconsin ice was concentrated in the areas adjacent to the mainland and Vancouver Island coasts.
9. Abrupt seaward thinning of the relatively short-lived late Wisconsin ice front in the vicinity of Queen Charlotte Sound suggests ice grounding may have extended in troughs no further than the shallower ledge on the northwestern slope of Moresby Trough and the mid-shelf moraines and ice front deposits in Mitchell's and Goose Island troughs, all of which lie at or near 200 m depth.
10. The height of bedrock platforms and/or volume of glacial debris deposited at any one site is largely what governed the discordant depths of shelf banks and not surf-scouring during a sea level minimum.
11. Intense tidal currents sweeping over banks after ice had retreated or was no longer grounded, but at a time prior to the sea attaining its present level, scoured bedrock and formed the well defined but very gently inclined valleys at the margins of Cook Bank. They also contributed to the formation of the heavy mineral deposits now found mainly at or near 100 m at the margins of Goose Island and Cook Bank.
12. The depth of the constructional (progradational) break in slope at the shelf edge has been controlled by relatively deep water oceanographic processes. At the time of the late Wisconsin glacial maximum, when the bulk of sediment was being discharged across the shelf, turbulence was probably high enough over the site of the present shelf edge to maintain at least sand and mud in suspension and permit continuous accumulation of material mainly below that depth.
13. The Coriolis effect and northward coastal water residual flow are directing the bulk of new sediment discharged into the Sound from inlets on the southern mainland coast towards the part of Goose Island Trough north of Sea Otter Shoals. Fine, suspended sediment also may be carried into the inner sound from the outer shelf by advective transport. Olive grey-green mud appears not to be accumulating at depths shallower than 140 m (except near the southeastern coast in the path of the plume of less saline water) and forms only a thin veneer (~30 cm) in Mitchell's Trough. For any given depth muds generally are finer on the more sheltered inner shelf and inlet mouths than on the outer shelf.
14. Estimated highest average Holocene sedimentation rates on the shelf (~2-3 mm/a) are found at isolated sites of fine sediment accumulation in northwestern Moresby Trough, at the bend in Mitchell's Trough and in Goose Island Trough north of the Sea Otter Shoals. The latter site may be the only location where sediments still are accumulating.

15. Smectite in olive grey surficial muds may have been formed as a consequence of present-day chemical "stripping" of chlorite, hornblende and mica derived from mainland rocks.
16. Most noncarbonate organic carbon is accumulating in deeper, more sheltered inshore parts of troughs and inlet mouths floored by smectite rich muddy sediments.
17. The relatively low concentrations of organic matter in subbottom, chlorite and illite rich greenish grey muds unconformably underlying surficial muds may reflect the low clay mineral content of constituent "glacial flour" and/or very rapid deposition which did not allow time for organic adsorption or development of a rich bottom fauna.
18. The most mobile glaciogene sediments in the sound are well sorted 2-3 ϕ sands. This population is being selectively washed from poorly sorted glacial debris on banks and gradually blanketing vast areas of bank surfaces. It is possible, however, that parts of bank surfaces are covered with relict outwash sands which were fairly well sorted when deposited. Sands will eventually be swept clear of bank tops by the combined effects of intense wave turbulence and tidal current transport.
19. Well sorted 2-3 ϕ sands are being swept (a) onto the northern flanks of Cook Bank, (b) towards the north and southwest of Goose Island Bank and (c) probably, at a slower, rate to the north and south of the shallower parts of Middle Bank.
20. The highest concentration of foraminifers within the sand plus mud fraction was found on Middle Bank, the deepest bank, in poorly sorted sediments scattered among boulders which may shelter the deposits somewhat from all but more extreme storm surges and tidal currents.
21. The highest concentrations of skeletal carbonate are found in a 10-15 km wide band extending along northern Vancouver Island and Scott Island Chain. A combination of fast currents, minimal terrigenous input and nutrient-rich upwelling slope waters probably have contributed to the development of this deposit.
22. This study has generated information:
 1. of direct relevance to local studies of groundfish/substrate associations and commercial groundfishing hazards.
 2. which may help design platforms suitable for local drilling conditions and which will help predict the behaviour of pollutants released in the course of any future hydrocarbon explorations.
 3. will help determine the economic potential of local sand, gravel, carbonate and heavy mineral deposits.

ACKNOWLEDGMENTS

J. Clague and C. Yorath critically reviewed earlier drafts of this report. T. Forbes helped proofread the final draft.

REFERENCES

- Anderson, J.B., Wheeler, R.B., and Schwarzer, R.B.
 1981: Sedimentologic and geochemical results of the Buccaneer oil/gas field study; in Proceedings 1981 Oil Spill Conference (Prevention, Behaviour, Control, Cleanup), March 2-5, 1981, Atlanta, Georgia, American Petroleum Institute, Environmental Protection Agency and United States Coast Guard, Washington, D.C., p. 721-724.
- Andrews, J.T. and Retherford, R.M.
 1978: A reconnaissance survey of Late Quaternary sea levels, Bella Bella / Bella Coola region, central British Columbia coast; Canadian Journal of Earth Sciences, v. 15, p. 341-350.
- Anonymous
 1980: Notes on placer mining in British Columbia; British Columbia Department of Mines and Petroleum Resources, Bulletin No. 21, 42 p.
- Armstrong, J.E.
 1966: Glacial studies, Kitimat - Terrace area; in Report of Activities, May to October 1963, Geological Survey of Canada, Paper 66-1, p. 50.
- Bader, R.G.
 1962: Some experimental studies with organic compounds and minerals - Graduate School of Oceanography, University of Rhode Island; Occasional Publication Number 1, 83 p.
- Baer, A.J.
 1967: Bella Coola and Laredo Sound map-areas, (93D, 103A) British Columbia; Geological Survey of Canada, Paper 66-25, 13 p.
- Baram, M.S., Rice, D., and Lee, W.
 1978: Marine mining of the continental shelf. Legal, technical and environmental considerations; Ballinger Publishing Company, Cambridge, Mass., 301 p.
- Barber, F.G.
 1957a: The effect of the prevailing winds on the inshore water masses of the Hecate Strait region, British Columbia; Journal of the Fisheries Research Board of Canada, 14, Number 6, p. 945-952.
 1957b: Observations of currents north of Triangle Island, British Columbia; Fisheries Research Board of Canada, Progress Reports of the Pacific Coast Stations, No. 108, p. 15-18.
 1958: Currents and water structure in Queen Charlotte Sound, British Columbia; Proceedings, Ninth Pacific Science Congress, 1957, Fisheries Research Board of Canada, No. 519, p. 196-199.
- Barnett, D.M. and Holdsworth, G.
 1974: Origin, morphology and chronology of sublacustrine moraines, Generator Lake, Baffin Island, Northwest Territories, Canada. Canadian Journal of Earth Sciences, v. 11, p. 380-408.
- Belderson, R.H. and Kenyon, N.H.
 1969: Direct illustration of one way sand transport by tidal currents; Journal of Sedimentary Petrology, v. 39, p. 1249-1250.

- Belderson, R.H., Kenyon, N.H., Stride, A.H., and Stubbs, A.R.
1972: Sonographs of the sea floor; Elsevier Scientific Publication Co. Amsterdam, 185 p.
- Bell, W.H.
1963: Surface current studies in the Hecate model; Fisheries Research Board of Canada, Manuscript Report, Number 159.
- Bevier, M.L., Armstrong, R.L., and Souther, J.G.
1979: Miocene peralkaline volcanism in west-central British Columbia - its temporal and plate tectonics settings; *Geology*, v. 7, p. 389-392.
- Biscaye, P.E.
1964: Distinction between kaolinite and chlorite in recent sediments by X-ray diffraction; *American Mineralogist*, v. 49, p. 1281-1289.
- Bjerkli, K. and Östmo-Saeter, J.S.
1973: Formation of glauconite in foraminiferal shells on the continental shelf off Norway; *Marine Geology*, v. 14, p. 169-179.
- Bradley, W.F. and Grim, R.E.
1961: Mica clay minerals; in *The X-ray identification and crystal structures of clay minerals*, ed. G. Brown; Mineralogical Society, London, p. 208-241.
- Brindley, G.W.
1961a: Experimental methods; in *The X-ray identification and crystal structures of clay minerals*, ed. G. Brown; Mineralogical Society, London, p. 1-50.
1961b: Chlorite minerals; in *The X-ray identification and crystal structures of clay minerals*, ed. G. Brown; Mineralogical Society, London, p. 242-296.
- Burst, J.F.
1958: Mineral heterogeneity in "glauconite" pellets; *American Mineralogist*, v. 43, p. 481-497.
- Canadian Hydrographic Service
1957: Co-tidal Charts - Pacific Coast of Canada (Preliminary Edition); Department of Mines and Technical Surveys, Canada.
- Carter, L.
1973: Surficial sediments of Barkley Sound and the adjacent continental shelf, west coast Vancouver Island, *Canadian Journal of Earth Sciences*, v. 10, p. 441-459.
- Chase, R.L., Tiffin, D.L., and Murray, J.W.
1975: The western Canadian continental margin; in *Canada's continental margins and offshore petroleum exploration*, eds. C.J. Yorath, E.R. Parker and D.J. Glass; Canadian Society of Petroleum Geologists, Memoir 4, p. 701-721.
- Chave, K.E.
1967: Recent carbonate sediments, an unconventional view; American Geological Institute, Council on Education in the Geological Sciences, *Journal of Geological Education*, v. 15, Short Reviews no. 7, p. 200-204.
- Clague, J.J.
1975: Late Quaternary sea level fluctuations, Pacific coast of Canada and adjacent areas; in *Report of Activities, Part C, Geological Survey of Canada, Paper 75-1C*, p. 17-21.
- Clague, J.J., Armstrong, J.E., and Mathews, W.H.
1980: Advance of the Late Wisconsin Cordilleran Ice Sheet in southern British Columbia since 22 000 yr. B.P.; *Quaternary Research*, v. 13, p. 322-326.
- Clague, J.J., Harper, J.R., Hebda, R.J., and Howes, D.E.
1982: Late Quaternary sea level and crustal movements, coastal British Columbia; *Canadian Journal of Earth Sciences*, v. 19, p. 597-618.
- Cockbain, A.E.
1963: Distribution of sediments on the continental shelf off the southern British Columbia coast; University of British Columbia, Institute of Oceanography, Manuscript Report, No. 15, 7 p.
- Crean, P.B.
1967: Physical oceanography of Dixon Entrance, British Columbia; Fisheries Research Board of Canada, Bulletin No. 156, 66 p.
- Culbert, R.R.
1971: A study of tectonic processes and certain geochemical abnormalities in the Coast Mountains of British Columbia; unpublished Ph.D. thesis, University of British Columbia, 229 p.
- Curry, J.R.
1969: History of continental shelves; in *The new concepts of continental margin sedimentation* (D.J. Stanley, convener) American Geological Institute, Washington, C.C., p. JC-6-1 to JC-6-18.
- Degens, E.T.
1965: *Geochemistry of sediments - a brief survey*; Prentice-Hall, New Jersey, 342 p.
- Dodimead, A.J.
1980: A general review of the oceanography of the Queen Charlotte Sound - Hecate Strait - Dixon Entrance region; Fisheries and Aquatic Sciences, Fisheries and Oceans Canada, Canadian Manuscript Report of Fisheries and Aquatic Sciences, No. 1574, 248 p.
- Dolan, R.
1970: Dune reddening along the outer banks of North Carolina; *Journal of Sedimentary Petrology*, v. 40, p. 765.
- Dreimanis, A.
1962: Quantitative gasometric determination of calcite and dolomite by using Chittick Apparatus; *Journal of Sedimentary Petrology*, v. 32, p. 520-529.
- Dubois, R.N.
1972: Inverse relation between foreshore slope and mean grain size as a function of the heavy mineral content; *Geological Association of America Bulletin*, v. 83, p. 871-876.
- Emery, K.L. and Niino, H.
1963: Sediments of the Gulf of Thailand and adjacent continental shelf; *Geological Society of America Bulletin*, v. 74, p. 541-554.
- Evans, O.F.
1939: Sorting and transportation of material in the swash and backwash; *Journal of Sedimentary Petrology*, v. 9, p. 28-31.
- Fillon, R.H.
1975: Deglaciation of the Labrador continental shelf; *Nature*, v. 253, p. 429-431.
1976: Hamilton Bank, Labrador shelf: post glacial sediment dynamics and paleo-oceanography; *Marine Geology*, v. 20, p. 7-25.
- Fladmark, K.R.
1974: A paleoecological model for Northwest Coast prehistory; unpublished Ph.D. thesis, University of Calgary, Calgary, Alberta, 319 p.

- Flint, R.F.
1937: Glacial and Pleistocene geology; Wiley and Sons, New York, 553 p.
- Folk, R.L.
1963: Petrology of sedimentary rocks; Hemphills, Texas, 159 p.
- Galehouse, J.S.
1971: Sedimentation analysis; in Procedures in sedimentary petrology, ed. R.E. Carver, John Wiley and Sons, New York, p. 69-94.
- Geological Society of America
1963: Rock Color Chart.
- Grant, A.C.
1972: The continental margin off Labrador and Eastern Newfoundland - morphology and geology; Canadian Journal of Earth Sciences, v. 9, p. 1394-1430.
- Hamilton, D., Sommerville, J.H., and Standord, P.N.
1980: Bottom currents and shelf sediments, southwest of Britain; Sedimentary Geology, v. 26, p. 115-138.
- Harms, J.C.
1969: Hydraulic significance of some sand ripples; Geological Society of America Bulletin, v. 80, p. 363-396.
- Hatcher, P.G. and Segar, D.A.
1976: Chemistry and continental margin sedimentation; in Marine sediment transport and environmental management, eds. D.J. Stanley and D.J.P. Swift, John Wiley & Sons, New York, p. 461-477.
- Hein, J.R., Allwardt, A.O., and Griggs, G.B.
1974: The occurrence of glauconite in Monterey Bay, California, diversity, origins, and sedimentary environmental significance; Journal of Sedimentary Petrology, v. 44, p. 562-571.
- Hein, J.R., Bouma, A.H., Hampton, M.A., and Ross, C.R.
1979: Clay mineralogy, fine grained sediment dispersal and inferred current patterns, lower Cook Inlet and Kodiak shelf, Alaska; Sedimentary Petrology, v. 24, p. 291-306.
- Holland, S.S.
1964: Landforms of British Columbia - a physiographic outline; Bulletin of the British Columbia Department of Mines and Petroleum Resources, No. 48, 138 p.
- Holmes, A.
1963: Principles of physical geology; The Ronald Press Co., New York, 1288 p.
- Holteidahl, H.
1958: Some remarks on geomorphology of continental shelves off Norway, Labrador, and southeast Alaska; Journal of Geology, v. 66, p. 461-571.
1960: Mountain, fiord, strand flat; geomorphology and general geology of parts of Western Norway; guide book "d" in Excursions in Norway, International Geological Congress, Oslo, Norway, 29 p.
1970: On the morphology of the West Greenland shelf with general remarks on the "marginal channel" problem; Marine Geology, v. 8, p. 155-172.
- Hoskin, C.M. and Nelson, R.V., Jr.
1969: Modern marine carbonate sediment, Alexander Archipelago, Alaska; Journal of Sedimentary Petrology, v. 39, p. 581-590.
- Houbolt, J.J.H.C.
1968: Recent sediments in the southern Bight of the North Sea; Geologie en Mijnbouw, v. 47, p. 247-273.
- Howes, D.E.
1981: Terrain inventory and geological hazards: northern Vancouver I.; Assessment and Planning Division Terrestrial Studies Branch, British Columbia Ministry of Environment, Bulletin 5, 105 p.
- Hughes, T.
1975: The West Antarctic ice sheet: instability, disintegration and initiation of ice ages; Reviews of Geophysics and Space Physics, v. 13, p. 502-526.
- Ingram, R.L.
1971: Sieve analysis; in Procedures in sedimentary petrology, ed. R.E. Carver, John Wiley and Sons, New York, p. 49-67.
- Inman, D.L.
1963: Sediments: properties, mechanics of sedimentation; in Submarine Geology, ed. F.P. Shepard, Harper and Row, New York, N.Y., 2nd ed.
- Jollymore, P.G.
1974: A medium range side-scan sonar for use in coastal waters: design criteria and operational experiences; in Proceedings I.E.E.E. Conference on Engineering in the Ocean Environment, Halifax, N.S., I.E.E. Inc., New York, 2, p. 108-114.
- Judd, J.B., Smith, W.C., and Pilkey, O.H.
1970: The environmental significance of iron-stained quartz grains on the southeastern United States Atlantic shelf; Marine Geology, v. 8, p. 355-362.
- Keen, C.E. and Hyndman, R.D.
1979: Geophysical review of the continental margins of eastern and western Canada; Canadian Journal of Earth Sciences, v. 16, p. 712-747.
- Keller, W.D.
1970: Environmental aspects of clay minerals; Journal of Sedimentary Petrology, v. 40, p. 788-854.
- Keller, G.H. and Richards, A.F.
1967: Sediments of the Malacca Strait, Southeast Asia; Journal of Sedimentary Petrology, v. 37, p. 102-127.
- Kenyon, N.H. and Stride, A.H.
1970: The tide swept continental shelf between the Shetland Isles and France; Sedimentology, v. 14, p. 159-173.
- King, L.H.
1969: Submarine end moraines and associated deposits on the Scotian Shelf; Geological Association of America Bulletin, v. 80, p. 83-96.
1976: Relict iceberg furrows on the Laurentian Channel and western Grand Banks; Canadian Journal of Earth Sciences, v. 13, p. 1082-1092.
- King, L.H., MacLean, B., and Drapeau, G.
1972: The Scotian shelf submarine end-moraine complex; Twenty-fourth International Geological Congress, Montreal; Marine Geology and Geophysics (Section 8), p. 237-249.

- Kodama, H. and Oinuma, K.
1963: Identification of kaolin minerals in the presence of chlorite by X-ray diffraction and infrared absorption spectra; in *Clays and clay minerals, Proceedings of the 13th National Conference on Clays and Clay Minerals*, Ottawa, 1962. Macmillan Co., New York, p. 236-249.
- Kulm, L.D., Robert, C.R., Harlett, J.C., Wendeck, R.H., Chambers, D.M., and Runge, E.J.
1975: Oregon continental shelf sedimentation: inter-relationships of facies distribution and sedimentary processes; *Journal of Geology*, v. 83, p. 145-175.
- Kunze, G.W., Knowles, L.J., and Kitano, Y.
1968: The distribution and mineralogy of clay minerals in the Taku Estuary of southeastern Alaska; *Marine Geology*, v. 6, p. 439-448.
- Ludwick, J.C.
1974: Tidal currents and zig-zag sand shoals in a wide estuary entrance; *Geological Society of America Bulletin*, v. 85, p. 717-726.
- Luternauer, J.L.
1972: Patterns of sedimentation in Queen Charlotte Sound, British Columbia; unpublished Ph.D. thesis, University of British Columbia, 203 p.
1976: Geofisheries research off the west coast of Canada; in *Report of Activities, Part C, Geological Survey of Canada, Paper 76-1C*, p. 157-159.
1978: Applications of side-scan sonar to geo-environmental research in the coastal waters of British Columbia; in *Current Research (Scientific and Technical Notes), Part B, Geological Survey of Canada, Paper 78-1B*, p. 181-186.
- MacEwan, D.M.C.
1961: Montmorillonite minerals; in *The X-ray identification and crystal structures of clay minerals*, ed. G. Brown; Mineralogical Society, London, p. 143-207.
- McBride, E.F.
1971: Mathematical treatment of size distribution data; in *Procedures in sedimentary petrology*, ed. R.E. Carver, John Wiley and Sons, New York, p. 109-127.
- McCave, I.N.
1972: Transport and escape of fine-grained sediment from shelf areas; in *Shelf sediment transport-process and pattern*, ed. D.J.P. Swift, D.B. Duane and O.H. Pilkey, Dowden, Hutchinson and Ross, Inc., Stroudsburg, Penn., p. 225-248.
- McGinnis, L.D.
1968: Glacial crustal bending; *Geological Society of America Bulletin*, v. 79, p. 769-776.
- McKinney, T.F., Stubblefield, W.L., and Swift, D.J.P.
1974: Large-scale current lineations on the Great Egg shoal retreat massif: investigations by side-scan sonar; *Marine Geology*, v. 17, p. 79-102.
- McLaren, P.
1982: The coastal geomorphology, sedimentology and processes of eastern Melville and western Byam Martin Islands, Canadian Arctic Archipelago; *Geological Survey of Canada, Bulletin 333*, 39 p.
- Mathews, W.H.
1958: Underwater gravel deposits of Dixon Entrance; University of British Columbia, Institute of Oceanography, Manuscript Report 7.
- May, J.P.
1973: Selective transport of heavy minerals by shoaling waves; *Sedimentology*, v. 20, p. 203-211.
- Miller, D.N. and Folk, R.L.
1955: Occurrence of detrital magnetite and ilmenite in red sediments; *American Association of Petroleum Geologists Bulletin*, 39, p. 338-345.
- Milliman, J.D.
1971: The role of calcium carbonate in continental shelf sedimentation; in *New concepts of continental margin sedimentation*; American Geological Institute Short Course Lecture Notes, Supplement, 20 p.
- Milne, W.G., Rogers, G.C., Riddihough, R.P., McMechan, G.A., and Hyndman, R.D.
1978: Seismicity of western Canada; *Canadian Journal of Earth Sciences*, v. 15, p. 1170-1193.
- Moore, T.C.
1973: Late Pleistocene-Holocene oceanographic changes in the northeastern Pacific, *Quaternary Research*, v. 3, p. 99-109.
- Muller, J. and Milliman, J.D.
1973: Relict carbonate - rich sediments on south-western Grand Bank, Newfoundland; *Canadian Journal of Earth Sciences*, v. 10, p. 1744-1750.
- Muller, J.E., Northcote, K.E., and Carlisle, D.
1974: Geology and mineral deposits of Alert Bay - Cape Scott Map Area, British Columbia; *Geological Survey of Canada, Paper 74-8*, 77 p.
- Muller, J.E.
1977: Evolution of the Pacific margin, Vancouver Island, and adjacent regions; *Canadian Journal of Earth Sciences*, v. 14, p. 2062-2085.
- Murray, J.W. and Mackintosh, E.E.
1968: Occurrence of interstratified glauconite-montmorillonite pellets, Queen Charlotte Sound; *Canadian Journal of Earth Sciences*, v. 5, p. 243-247.
- Naidu, A.S., Burrell, D.C., Hood, D.W., and Dygas, J.A.
1975: Texture clay mineralogy, and chemistry of bottom sediments, West Beaufort Sea, Arctic Ocean; in *Contributions to the geology of the Bering Sea basin and adjacent regions*, ed. R.B. Forbes, The Geological Society of America, Special Paper 151, p. 49-58.
- National Research Council
1975: Mining in the outer continental shelf and in the deep ocean; Panel on Operational Safety in Marine Mining, Marine Board, Assembly of Engineering, National Academy of Sciences, Washington, D.C.
- Nayudu, Y.R. and Enbysk, B.J.
1964: Bio-lithology of northeast Pacific surface sediments; *Marine Geology*, v. 2, p. 310-342.
- O'Brien, N.R. and Burrell, D.C.
1970: Mineralogy and distribution of clay size sediment in Glacier Bay, Alaska; *Journal of Sedimentary Petrology*, v. 40, p. 650-655.
- Off, T.
1963: Rhythmic linear sand bodies caused by tidal currents; *American Association of Petroleum Geologists Bulletin*, v. 47, p. 324-341.
- Ollier, C.
1969: *Weathering*; Oliver and Boyd, Edinburgh, 304 p.

- Pacific Oceanographic Group
1955a: Physical and chemical data record, Hecate Project, 1954; Joint Committee on Oceanography Manuscript Report, 99 p.
1955b: Physical and chemical data record. Hecate Project with Appendix 1 - current observations, 1955; Joint Committee on Oceanography, Manuscript Report, 107 p.
- Peacock, M.A.
1935: Fiord-land of British Columbia; Bulletin of the Geological Society of America, v. 46, p. 633-696.
- Pickard, G.L. and Giovando, L.F.
1960: Some observations of turbidity in British Columbia inlets; Limnology and Oceanography, v. 5, p. 162-170.
- Rao, C.B.
1957: Beach erosion and concentration of heavy mineral sands; Journal of Sedimentary Petrology, v. 27, p. 143-147.
- Riddihough, R.P.
1977: A model for recent plate interactions off Canada's west coast; Canadian Journal of Earth Sciences, v. 14, p. 384-396.
1982: Contemporary vertical movements and tectonics on Canada's west coast: a discussion; Tectonophysics, v. 86, p. 319-341.
- Roddick, J.A., Baer, A.J., and Hutchinson, W.W.
1966: Coast mountains project; in Report of Activities, May to October 1965, Geological Survey of Canada, Paper 66-1, p. 80-85.
- Roddick, J.A., Mulier, J.E., and Okulitch, A.V.
1976: Fraser River, British Columbia - Washington; Geological Survey of Canada, Energy, Mines and Resources Canada, Map 1386A, Sheet 92.
- Rogers, G.C.
1982: Revised seismicity and revised fault plane solutions for the Queen Charlotte Islands Region; Earth Physics Branch, Open File No. 82-23, 50 p.
- Schafer, C.T. and Prakash, A.
1968: Current transport and deposition of Foraminiferal tests, plankton organisms and lithogenic particles in Bedford Basin, Nova Scotia; Maritime Sediments, v. 4, p. 100-103.
- Shouldice, D.H.
1971: Geology of the western Canadian continental shelf; Canadian Petroleum Geology Bulletin, v. 19, p. 405-436.
1973: Western Canadian continental shelf; in Future Petroleum Provinces of Canada - Their Geology and Potential, ed. R.G. McGrossan, Canadian Society of Petroleum Geologists, Memoir 1, p. 7-35.
- Southard, J.B. and Stanley, D.J.
1976: Shelf-break processes and sedimentation; in Marine Sediment Transport and Environmental Management, eds. D.J. Stanley and D.J.P. Swift, John Wiley & Sons, New York, p. 351-377.
- Souther, J.G.
1966: Cordillera volcanic study; in Report of Activities, May to October, 195, Geological Survey of Canada, Paper 66-1, p. 87-89.
- Stacey, R.A.
1975: Structure of the Queen Charlotte Basin; in Canada's continental margins and offshore petroleum exploration, ed. C.J. Yorath, E.R. Parker and D.J. Glass, Canadian Society of Petroleum Geologists, Memoir 4, p. 701-721.
- Stanley, D.J.
1968: Reworking of glacial sediments in the North West Arm, a fiord-like inlet on the southeast coast of Nova Scotia; Journal of Sedimentary Petrology, v. 38, p. 1224-1241.
- Sutherland Brown, A.
1966: Tectonic history of the Insular Belt of British Columbia; in Tectonic history and mineral deposits of the western Cordillera, Canadian Institute of Mining and Metallurgy, Montreal, p. 83-100.
1968: Geology of the Queen Charlotte Islands, British Columbia; British Columbia Department of Mines and Petroleum Resources, Bulletin 54, 226 p.
- Swift, J.D.P.
1969: Outer shelf sedimentation: processes and products; in The new concepts of continental shelf sedimentation, convenor D.J. Stanley, AGI Short Course Lecture Notes; American Geological Institute, Washington, p. DS-5-1 to DS-5-26.
- Swift, D.J.P. and Boehmer, W.R.
1972: Brown and grey sands on the Virginia shelf: color as a function of grain size; Geological Society of America Bulletin, v. 83, p. 877-884.
- Swift, D.J.P. and Freeland, G.L.
1978: Current lineations and sand waves on the inner shelf, middle Atlantic Bight of North America; Journal of Sedimentary Petrology, v. 48, p. 1257-1266.
- Swift, D.J.P., Nelson, T., McHone, J., Holliday, B., Palmer, H., and Shideler, G.
1978: Holocene evolution of the inner shelf of southern Virginia; Journal of Sedimentary Petrology, v. 47, p. 1454-1474.
- Thomas, R.H. and Bentley, C.R.
1978: A model for Holocene retreat of the West Antarctic ice sheet; Quaternary Research, v. 10, p. 150-170.
- Thomas, R.L.
1969: A note on the relationship of grain size, clay content, quartz and organic carbon in some Lake Erie and Lake Ontario sediments; Journal of Sedimentary Petrology, v. 39, p. 803-809.
- Thomson, R.E.
1981: Oceanography of the British Columbia coast; Canadian Special Publication of Fisheries and Aquatic Sciences 56, Fisheries and Oceans Canada, 56, 291 p.
- Tully, J.P.
1952: Climate in the coastal seas of British Columbia; Fisheries Research Board of Canada, Progress Reports of the Pacific Coast Stations, No. 90, p. 16-20.
- University of British Columbia Institute of Oceanography
1963: Sediment grain size analyses; Data Report, No. 22, 6 p.

Van der Meer, W.J., Fillon, R.H., and Monshaw, D.

- 1976: Hamilton Bank, Labrador margin: origin and evolution of a glaciated shelf; Department of the Environment, Marine Sciences Paper 14 and Geological Survey of Canada, Paper 75-40, 31 p.

Walker, G.F.

- 1961: Vermiculite minerals; in The X-ray identification and crystal structures of clay minerals; ed. G. Brown, Mineralogical Society, London, p. 297-324.

Warshaw, C.M., Rosenberg, P.E., and Ruston, R.

- 1960: Changes effected in layer silicates by heating below 550°C.; Clay Minerals Bulletin, 4, p. 113-126.

Weaver, C.E.

- 1958: The effect and geologic significance of potassium "fixation" by expandable clay minerals derived from muscovite, biotite, chlorite and volcanic mineral; American Mineralogist, v. 43, p. 839-861.

Westrheim, S.J.

- 1974: Echo-sounder and trawl survey of Queen Charlotte Sound and southern Hecate Strait, 1972-73; Fisheries Research Board of Canada, Manuscript Report, No. 1307.

Wiese, W.

- 1969: Studies on properties, distribution, and heavy mineral contents of sediments in northern Queen Charlotte Sound; unpublished B.Sc. thesis, Department of Geological Sciences, University of British Columbia, Department of Geological Sciences, 78 p.

Yorath, C.J., Bornhold, B.D., and Thomson, R.E.

- 1979: Oscillation ripples on the northeast Pacific continental shelf; Marine Geology, v. 31, p. 45-58.

Yorath, C.J. and Chase, R.L.

- 1981: Tectonic history of the Queen Charlotte Islands and adjacent areas - a model; Canadian Journal of Earth Sciences, v. 18, p. 1717-1739.

APPENDIX
Primary characteristics of samples

EXAMPLE:

QCS# 254 52°13' 129°00' 133
1.492 2.225 -0.488 12.360 79.570 8.070 0.5 16

WHICH REPRESENTS:

SAMPLE NO. LOCATION DEPTH IN METRES

MEAN PHI/STANDARD/SKEWNESS
DEVIATION PHI

WT%
GRAVEL

WT%
SAND

WT%
MUD

WT%
NON-CO₂
ORGANIC
CARBON

WT%
CO₂

QCS# 2	51°08'	127°45'	73	0.0	97.023	2.977	0.3	1	QCS# 29	50°54'	128°40'	88	89.100	10.809	0.090	0.0	0
3.294	0.462	0.489							-3.941	2.121	0.925						
QCS# 3	51°05'	127°51'	120	1.773	72.859	25.367	0.8	2	QCS# 30	50°52'	128°45'	55	93.423	6.560	0.017	0.0	0
3.484	1.994	0.525							-4.784	1.707	1.331						
QCS# 4	51°03'	127°55'	167	0.0	86.242	13.758	0.6	1	QCS# 32	50°53'	128°54'	60	(trace of gravel)				
3.352	1.815	1.452							QCS# 33	50°55'	128°09'	64					
QCS# 5	51°01'	128°01'	146	0.0	94.880	5.120	0.4	2	-2.762	2.409	0.611	79.933	19.675	0.392	0.2	21	
2.922	0.715	0.454							QCS# 34	50°57'	128°44'	82					
QCS# 6	50°59'	128°06'	51	88.428	11.335	0.238	0.0	0	2.200	0.484	-0.493	0.0	99.937	0.063	0.0	2	
-3.160	1.980	0.914							QCS# 35	51°00'	128°37'	66	6.404	93.585	0.011	0.1	0
QCS# 7	50°57'	128°11'	55	100.00	0.0	0.0	0.0	0	0.603	1.208	-0.879						
-6.048	0.594	0.112							QCS# 36	51°02'	128°32'	95	0.0	99.791	0.209	0.1	1
QCS# 8	50°54'	128°17'	37	100.000	0.0	0.0	0.0	0	2.776	0.384	0.201						
-5.416	0.455	0.453							QCS# 37	51°04'	128°27'	122	0.0	97.426	2.574	0.1	1
QCS# 9	50°52'	128°22'	44	16.862	83.090	0.049	0.0	20	2.800	0.678	0.067						
-0.028	1.260	-0.430							QCS# 38	51°06'	128°21'	175	0.0	74.398	25.602	0.6	1
QCS# 11	50°51'	128°37'	49	54.897	43.721	1.382	0.0	20	3.636	1.461	1.365						
-1.081	2.098	0.244							QCS# 39	51°09'	128°16'	165	1.116	85.236	13.648	0.5	1
QCS# 10	50°50'	128°28'	53	(trace of gravel)					3.035	2.232	1.076						
QCS# 12	50°54'	128°29'	64	6.029	93.810	0.161	0.1	16	QCS# 40	51°10'	128°12'	99	0.0	96.691	3.309	0.0	0
1.884	1.405	-1.084							2.107	0.851	0.537						
QCS# 15	51°00'	128°15'	80	0.0	99.919	0.081	0.0	1	QCS# 41	51°13'	128°17'	95	0.390	98.776	0.835	0.0	0
2.362	0.529	-0.087							2.257	0.636	-0.599						
QCS# 16	51°02'	128°10'	163	0.0	80.231	19.770	0.7	3	QCS# 42	51°12'	128°21'	146	0.0	91.194	8.806	0.5	2
3.407	1.641	1.329							3.235	0.674	0.621						
QCS# 17	51°04'	128°05'	153	0.0	81.132	18.868	0.7	2	QCS# 43	51°10'	128°25'	192	0.0	73.849	26.151	0.7	1
4.046	2.049	1.175							3.907	1.176	2.195						
QCS# 18	51°06'	128°00'	100	60.132	25.060	14.808	1.0	8	QCS# 44	51°08'	128°32'	161	0.0	89.570	10.430	0.4	0
-1.049	4.492	0.503							3.489	0.853	2.962						
QCS# 19	51°09'	127°54'	119	0.0	47.740	52.260	0.9	2	QCS# 45	51°06'	128°36'	119	0.0	97.206	2.794	0.2	1
5.028	2.410	0.932							3.129	0.484	0.702						
QCS# 20	51°11'	127°49'	49	49.458	49.273	1.269	0.0	0	QCS# 46	51°04'	128°41'	64	0.0	99.983	0.017	0.0	0
-0.588	3.299	0.008							1.347	0.652	0.135						
QCS# 21	51°12'	127°57'	123	0.0	64.798	35.202	0.7	2	QCS# 47	51°01'	128°47'	55	(trace of gravel)				
4.304	2.037	1.175							QCS# 48	50°59'	128°52'	68	92.204	7.762	0.034	0.0	0
QCS# 22	51°10'	128°03'	68	74.920	23.188	1.892	0.0	0	-3.272	1.684	0.840						
-2.952	3.043	0.581							QCS# 49	50°57'	128°58'	68	0.0	99.939	0.061	0.1	0
QCS# 23	51°08'	128°08'	128	72.040	15.413	12.547	0.0	0	2.222	0.431	-0.007						
-1.795	4.539	0.736							QCS# 50	50°55'	129°04'	68	31.425	68.550	0.024	0.0	0
QCS# 24	51°06'	128°13'	166	0.0	88.060	11.941	0.5	1	-0.238	1.726	-0.276						
2.961	1.435	1.613							QCS# 51	50°52'	129°11'	62	100.000	0.0	0.0	0.0	0
QCS# 25	51°04'	128°19'	164	0.0	72.318	27.682	0.7	2	-6.297	0.630	0.712						
4.119	1.630	1.625							QCS# 55	50°51'	129°22'	168	0.0	94.354	5.646	0.5	1
QCS# 26	51°02'	128°24'	99	30.914	68.378	0.708	0.0	1	2.855	0.640	1.004						
0.165	2.988	-0.381							QCS# 56	50°53'	129°18'	123	26.401	70.336	3.263	0.7	26
QCS# 27	50°59'	128°29'	91	0.0	99.462	0.538	0.1	1	0.392	2.173	-0.157						
2.674	0.530	-0.202							QCS# 57	50°55'	129°12'	88	(trace of gravel)				
QCS# 28	50°56'	128°34'	55	98.777	1.202	0.021	0.0	0									
-5.166	1.050	1.908															

QCS# 58 2.578	50°58' 0.428	129°7' 0.707	100 0.0	99.213	0.787	0.2	2	QCS# 90 4.760	51°26' 1.745	128°34' 1.163	207 0.0	33.661	66.339	0.3	2
QCS# 59 2.428	51°06' 0.460	129°1' -0.196	86 0.0	99.849	0.151	0.1	1	QCS# 91 6.372	51°28' 1.886	128°29' 0.404	194 0.0	15.305	84.695	0.9	3
QCS# 60 1.892	51°2' 0.548	128°56' 0.272	84 0.0	99.779	0.221	0.0	0	QCS# 92 6.224	51°29' 2.912	128°21' 0.404	157 0.0	16.429	83.571	0.3	3
QCS# 61 -1.036	51°5' 2.521	128°50' -0.131	80 47.803	52.017	0.180	0.1	6	QCS# 93 4.128	51°32' 1.878	128°18' 1.392	154 0.0	68.909	31.090	0.8	8
QCS# 62 0.792	51°8' 0.897	128°45' -0.124	73 2.319	97.669	0.012	0.0	0	QCS# 94 2.889	51°34' 0.351	128°13' 0.604	100 0.0	99.509	0.491	0.0	1
QCS# 63 2.547	51°9' 0.509	128°40' 0.160	122 0.0	99.713	0.287	0.0	0	QCS# 97 5.487	51°44' 2.469	128°23' 0.756	141 0.0	26.502	73.499	1.3	3
QCS# 64 3.373	51°11' 1.091	128°35' 2.641	170 0.0	92.674	7.326	0.4	1	QCS# 98 4.548	51°47' 2.332	128°26' 0.917	155 0.0	63.961	36.039	1.7	3
QCS# 65 3.871	51°14' 1.236	128°29' 1.993	192 0.0	74.336	25.663	1.0	2	QCS# 99 3.702	51°46' 1.095	128°32' 2.196	146 0.0	82.910	17.090	0.5	1
QCS# 66 -4.963	51°16' 1.095	128°24' 2.030	111 98.510	1.440	0.050	0.0	0	QCS# 100 1.881	51°42' 0.495	128°37' -0.034	92 0.0	99.982	0.018	0.0	1
QCS# 68 0.644	51°18' 3.375	128°29' 0.649	133 40.910	43.839	15.251	0.7	9	QCS# 101 2.003	51°41' 0.431	128°42' -0.325	37 0.0	99.985	0.015	0.0	1
QCS# 69 -1.024	51°16' 2.979	128°32' -0.031	183 42.588	54.022	3.390	0.3	14	QCS# 102 1.611	51°38' 0.660	128°47' -1.110	40 1.598	98.396	0.006	0.0	3
QCS# 71 2.927	51°12' 0.459	128°42' 0.676	174 0.0	98.516	1.484	0.3	0	QCS# 103 -1.495	51°36' 1.696	128°52' 0.356	46 68.033	31.937	0.031	0.0	4
QCS# 72 1.193	51°10' 1.071	128°48' -0.229	110 2.981	96.958	0.062	0.0	0	QCS# 104 -2.298	51°34' 1.860	128°58' 0.552	46 76.606	23.394	0.0	0.0	0
QCS# 73 2.306	51°8' 0.735	128°53' -0.550	100 0.0	99.780	0.220	0.1	0	QCS# 115 3.638	51°00' 1.350	129°23' 2.043	183 0.0	89.250	10.750	0.4	3
QCS# 74 2.801	51°5' 0.391	128°59' 0.119	102 0.0	99.852	0.148	0.1	1	QCS# 116 3.723	51°02' 1.407	129°18' 1.967	174 0.0	83.527	16.473	0.6	4
QCS# 75 1.498	51°3' 2.696	129°5' -0.560	124 19.531	75.088	5.381	0.4	3	QCS# 117 4.054	51°04' 1.695	129°13' 1.744	155 0.0	76.276	23.724	0.7	4
QCS# 76 3.586	51°1' 1.053	129°10' 2.154	145 0.0	88.092	11.908	0.7	4	QCS# 118 2.539	51°06' 1.053	129°08' 0.098	137 0.0	96.104	3.896	0.5	5
QCS# 77 3.107	50°58' 1.258	129°16' 2.488	157 0.0	95.069	4.931	0.7	3	QCS# 119 1.573	51°09' 1.209	129°02' -0.175	137 3.348	95.294	1.359	0.0	1
QCS# 78 3.477	50°56' 1.759	129°21' 1.553	173 0.0	86.973	13.027	0.6	9	QCS# 120 -3.453	51°12' 2.250	128°58' 0.941	137 85.639	14.064	0.297	0.2	28
QCS# 79 3.686	50°54' 1.515	129°28' 1.624	192 0.0	81.852	18.149	0.6	9	QCS# 121 0.578	51°14' 1.978	128°52' -0.259	146 32.261	67.716	0.023	0.0	0
QCS# 80 3.529	51°4' 1.417	129°29' 1.674	280 0.359	85.326	14.315	0.6	4	QCS# 124 -3.411	51°18' 3.298	128°41' 0.407	183 72.190	26.817	0.993	0.3	2
QCS# 81 -1.400	51°7' 3.630	129°24' 0.280	265 60.994	35.120	3.885	1.0	9	QCS# 125 2.352	51°20' 1.010	128°35' -0.172	183 0.0	95.102	4.898	0.2	1
QCS# 82 3.389	51°9' 1.892	129°18' 1.180	238 0.0	84.033	15.967	0.8	9	QCS# 126 4.025	51°23' 1.702	128°31' 1.589	174 0.0	73.927	26.073	0.6	1
QCS# 83 1.8883	51°11' 3.432	129°12' -0.061	223 19.975	67.289	12.736	0.6	3	QCS# 127 4.777	51°25' 2.291	128°25' 1.019	164 0.0	53.465	46.535	1.2	3
QCS# 84 2.714	51°13' 0.964	129°7' 0.266	230 0.0	91.942	8.058	0.5	1	QCS# 129 3.458	51°24' 1.193	128°16' 1.944	128 0.0	84.722	15.278	0.4	3
QCS# 85 2.479	51°14' 0.786	129°1' 0.872	205 0.0	94.658	5.342	0.3	1	QCS# 131 2.716	51°20' 0.410	128°08' 0.475	137 0.0	99.280	0.720	0.8	5
QCS# 86 3.491	51°17' 1.461	128°56' 2.033	223 0.0	88.445	11.555	0.4	1	QCS# 132 4.067	51°26' 1.461	128°12' 1.505	137 0.0	61.556	38.445	1.4	3
QCS# 87 3.956	51°19' 2.178	128°49' 1.084	208 0.907	80.068	19.024	0.6	2	QCS# 133 1.936	51°29' 1.042	128°15' 0.149	100 0.0	97.485	2.515	0.2	3
QCS# 88 2.276	51°21' 2.696	128°45' 0.086	220 12.640	74.256	13.104	05.	1	QCS# 134 2.279	51°30' 0.588	128°12' -0.596	91 0.967	97.941	1.092	0.1	0
QCS# 89 4.407	51°23' 1.884	128°39' 1.345	214 0.0	61.598	38.402	0.8	1	QCS# 137 5.375	51°38' 2.448	128°18' 0.793	146 0.0	50.722	69.278	1.3	2

QCS# 138 6.074	51033' 2.708	128024' 0.528	174 0.0	15.752	84.248	2.3	3	QCS# 177 -3.294	51033' 1.689	128050' 0.892	46 91.630	8.352	0.018	0.0	0
QCS# 139 5.891	51032' 2.590	128028' 0.583	183 0.0	22.172	77.828	2.4	3	QCS# 178 -3.617	51031' 2.803	128056' 0.394	46 73.041	26.943	0.016	0.1	1
QCS# 140 5.439	51030' 2.522	128034' 0.723	183 0.0	34.377	65.623	1.4	2	QCS# 180 2.438	51025' 0.544	12907' -0.040	82 0.0	99.890	0.110	0.0	0
QCS# 141 4.320	51028' 2.032	128040' 1.240	183 0.0	67.058	32.942	1.1	2	QCS# 181 2.242	51025' 0.641	129012' 0.073	110 0.0	99.753	0.247	0.1	0
QCS# 142 4.083	51027' 1.412	128044' 1.720	183 0.0	66.739	33.261	0.8	2	QCS# 182 2.422	51023' 0.374	129018' 0.047	146 0.0	99.961	0.039	0.1	0
QCS# 143 4.626	51025' 2.123	128049' 1.169	200 0.0	56.467	43.533	0.9	2	QCS# 183 3.369	51020' 0.565	129023' 0.751	192 0.0	92.653	7.347	0.4	1
QCS# 144 4.612	51023' 2.26	128055' 1.012	220 0.0	57.695	42.305	0.8	1	QCS# 184 2.012	51019' 3.919	129029' -0.131	220 20.020	62.674	17.306	0.8	5
QCS# 145 2.484	51021' 0.804	12900' 0.648	230 0.0	94.605	5.395	0.2	1	QCS# 185 -3.824	51017' 1.900	129034' 1.113	238 90.393	9.531	0.076	0.0	0
QCS# 146 3.641	51019' 1.597	12905' 1.778	256 0.0	85.531	14.469	0.5	1	QCS# 187 3.648	51012' 2.023	129043' 1.180	457 0.0	81.260	18.740	0.2	3
QCS# 147 2.675	51016' 0.900	12909' 0.649	256 0.0	91.236	8.764	0.3	1	QCS# 225 2.417	51018' 1.113	127055' 0.347	119 0.0	90.366	9.634	0.5	3
QCS# 148 3.989	51014' 1.330	129015' 1.711	284 0.0	66.305	33.695	0.7	3	QCS# 226 -0.354	51024' 3.181	127053' -0.080	111 42.475	53.676	3.849	0.3	1
QCS# 149 0.300	51012' 4.169	129019' -0.143	293 37.723	49.586	12.691	0.6	5	QCS# 227 4.909	51022' 2.612	127058' 0.820	440 0.0	54.910	44.560	1.3	4
QCS# 151 3.674	5108' 1.198	129031' 2.108	284 0.0	86.616	13.384	0.5	4	QCS# 228 2.631	51025' 1.043	12802' 0.299	110 0.0	90.044	9.956	0.5	1
QCS# 152 3.978	5105' 1.563	129037' 1.1593	450 0.0	79.981	20.019	0.5	3	QCS# 229 6.644	51050' 2.608	128020' 0.381	154 0.0	9.322	90.677	3.6	5
QCS# 154 -3.273	5106' 2.376	129046' 0.843	420 (trace of gravel) 274	84.611	13.990	1.398	14	QCS# 230 -0.269	51052' 0.660	128030' 0.114	71 9.123	90.860	0.017	0.0	0
QCS# 155 -3.273	5109' 2.376	129040' 0.843	274 84.611	13.990	1.398	0.0	14	QCS# 231 -3.218	51050' 3.707	128035' 0.636	137 75.536	20.417	4.048	0.6	4
QCS# 156 51010'	129036'	274 (trace of gravel)	274					QCS# 232 -1.794	51055' 3.270	128033' 0.108	88 59.213	40.500	0.287	0.2	7
QCS# 158 4.873	51015' 1.983	129025' 1.135	293 0.0	36.057	63.944	1.4	4	QCS# 233 -1.449	5200' 1.293	128031' 0.680	64 72.514	27.151	0.334	0.0	0
QCS# 162 4.373	51023' 1.825	12903' 1.431	200 0.0	63.008	36.992	0.8	2	QCS# 234 0.974	5204' 1.443	128034' -1.153	71 8.710	90.957	0.333	0.5	30
QCS# 163 2.980	51025' 1.344	128053' 1.818	170 0.0	90.617	9.383	0.3	1	QCS# 235 4.866	5207' 2.031	128038' 1.012	214 0.0	39.631	60.369	0.9	4
QCS# 164 3.148	51027' 0.501	128052' 0.846	155 0.0	96.280	3.720	0.0	0	QCS# 236 8.164	52013' 2.710	128036' -0.087	238 0.0	5.673	94.327	3.3	7
QCS# 165 1.888	51029' 0.554	128046' 0.047	73 0.0	99.961	0.039	0.0	0	QCS# 237 7.449	52011' 2.821	128042' 0.088	238 0.0	10.973	89.026	3.2	6
QCS# 166 2.045	51032' 0.415	128041' -0.257	40 0.0	99.972	0.028	0.0	4	QCS# 238 4.398	52015' 2.009	128046' 1.289	192 0.0	65.202	34.797	1.6	8
QCS# 167 3.678	51034' 1.074	128035' 2.137	146 0.0	82.910	17.090	0.1	2	QCS# 239 4.617	52018' 2.440	128049' 0.810	205 0.0	63.176	36.824	1.6	10
QCS# 168 1.821	51037' 3.944	128030' -0.064	150 24.015	50.553	25.431	0.8	2	QCS# 241 -5.675	52019' 1.046	128058' 0.186	55 100.000	0.0	0.0	0.0	0
QCS# 169 4.177	51039' 1.363	128025' 1.699	146 0.0	56.655	43.345	0.7	1	QCS# 242 -5.595	52016' 0.583	128054' 1.887	92 100.000	0.0	0.0	0.0	0
QCS# 170 3.331	51042' 1.168	128021' 1.947	137 0.0	87.576	12.424	0.5	1	QCS# 243 4.546	52012' 3.587	128051' 0.119	210 6.594	48.930	44.476	1.3	5
QCS# 171 4.726	51045' 2.116	128018' 0.919	146 0.0	43.126	56.875	1.5	3	QCS# 244 6.372	5209' 2.722	128048' 0.586	214 0.0	17.799	82.202	2.7	5
QCS# 173 4.232	51041' 1.533	128030' 1.713	146 0.0	65.689	34.310	0.6	2	QCS# 245 4.627	5205' 1.856	128043' 1.275	183 0.0	49.067	50.933	1.0	3
QCS# 175 2.178	51050' 0.345	128040' 0.254	36 0.0	99.845	0.155	0.0	2	QCS# 246 4.139	5202' 1.226	128041' 1.783	164 0.0	59.913	40.087	0.6	3

QCS# 247 4.449	51058' 1.987	128027' 1.114	155 0.0	61.939	38.061	0.6	2	QCS# 277 -3.098	52016' 3.196	129017' 0.993	146 81.695	13.914	4.391	0.0	0
QCS# 248 3.538	51053' 1.112	128042' 2.284	139 0.0	87.188	12.812	0.3	2	QCS# 278 -3.941	52014' 2.361	129022' 0.937	153 90.851	7.576	1.573	0.0	0
QCS# 249 4.207	51057' 1.444	128042' 1.741	148 0.0	58.967	41.033	0.4	4	QCS# 279 -1.758	52010' 2.722	129019' 0.356	159 65.297	32.084	2.619	0.7	9
QCS# 250 4.328	52001' 1.466	128046' 1.601	164 0.0	54.884	45.116	0.3	4	QCS# 280 2.268	5207' 1.184	129015' 0.281	153 0.0	91.219	8.781	0.4	4
QCS# 251 4.289	5204' 1.891	128049' 1.883	168 0.0	74.103	25.897	0.4	1	QCS# 281 3.517	5203' 1.336	129012' 1.766	168 0.0	79.806	20.194	0.6	4
QCS# 252 3.355	5206' 3.086	128053' 0.332	189 8.959	64.603	26.438	1.3	3	QCS# 282 -2.704	5200' 3.529	129009' 0.401	164 68.607	27.816	3.577	0.2	4
QCS# 253 -0.602	52010' 3.227	128057' 0.022	170 46.839	46.843	6.318	0.4	10	QCS# 284 3.381	51053' 0.630	129011' 0.470	111 0.0	91.492	8.508	0.0	1
QCS# 254 1.492	52013' 2.225	12900' -0.488	133 12.360	79.570	8.070	0.5	16	QCS# 285 -1.563	51049' 3.477	129058' 0.154	84 57.409	41.643	0.947	0.1	2
QCS# 255 4.074	52017' 3.717	12903' 0.383	153 5.036	62.995	31.969	0.7	7	QCS# 286 1.630	51046' 0.848	128055' -0.664	49 1.465	98.535	0.0	0.0	2
QCS# 256 1.666	52020' 3.673	12907' 0.039	141 24.314	61.949	13.736	0.6	6	QCS# 287 2.756	51043' 0.464	128051' -0.361	71 0.0	99.622	0.378	0.1	4
QCS# 257 3.978	52020' 1.955	129011' 1.151	157 0.0	73.770	26.229	0.7	8	QCS# 288 2.531	51038' 0.710	128052' -1.494	38 (camera station)				
QCS# 258 6.297	52015' 3.195	12908' 0.332	166 0.0	32.655	67.345	1.3	6	QCS# 289 2.564	51042' 0.722	128057' -0.241	62 0.0	98.547	0.234	0.0	4
QCS# 259 -1.459	52011' 2.849	12904' 0.110	113 56.128	42.670	1.201	0.1	13	QCS# 290 3.726	51045' 0.992	12901' 2.717	81 0.0	99.066	0.934	0.1	10
QCS# 260 -0.755	52013' 2.948	12902' -0.103	128 43.927	54.018	2.055	0.2	9	QCS# 292 2.373	51051' 0.383	12907' 0.732	117 0.0	85.655	14.345	0.2	
QCS# 261 4.001	5205' 1.381	128057' 1.971	157 0.0	75.322	24.678	1.3	2	QCS# 293 5.077	51054' 2.399	129011' 0.963	137 0.0	99.621	0.379	0.0	0
QCS# 262 -1.524	5201' 2.666	128055' 0.242	183 63.191	35.615	1.194	0.5	10	QCS# 294 2.565	51058' 1.143	129014' 0.928	175 0.0	37.108	62.892	0.8	0
QCS# 263 0.489	51056' 1.787	128051' -0.430	73 23.763	76.137	0.100	0.0	1	QCS# 295 0.387	5201' 2.399	129018' 0.928	186 0.0	89.038	10.962	0.1	3
QCS# 264 3.074	51054' 0.406	128048' 0.551	95 0.0	98.885	1.115	0.1	1	QCS# 296 5.138	5205' 2.335	129022' 0.309	186 42.968	42.157	14.875	0.5	6
QCS# 265 3.128	51051' 0.424	128044' 0.415	75 0.0	98.585	1.415	0.0	2	QCS# 297 6.453	5208' 2.983	129024' 0.909	200 0.0	36.286	63.714	1.0	5
QCS# 266 2.852	51047' 0.509	128040' 0.367	77 0.0	98.673	1.327	0.0	3	QCS# 298 5.143	52011' 2.244	129030' 0.266	204 0.0	20.943	79.056	1.2	6
QCS# 267 2.525	51043' 0.723	128043' -1.089	37 (camera station)					QCS# 299 0.607	52014' 2.244	129033' 0.830	214 0.0	50.270	49.730	1.2	4
QCS# 268 2.525	51045' 0.723	128047' -1.089	55 0.820	98.792	0.388	0.0	3	QCS# 300 4.613	52012' 1.853	129038' 1.346	218 0.0	47.129	52.871	0.6	4
QCS# 269 2.492	51048' 0.607	128049' -0.480	64 0.0	99.751	0.249	0.0	1	QCS# 301 4.957	5209' 2.974	129034' 0.320	216 2.498	36.639	60.863	0.9	6
QCS# 270 1.391	51052' 1.948	128053' -0.654	62 13.980	83.716	2.304	0.0	0	QCS# 311 3.060	51033' 0.465	12904' 0.549	50 (camera station)				
QCS# 271 2.406	51055' 0.488	128056' -0.236	80 0.0	99.646	0.354	0.1	1	QCS# 312 2.740	51043' 0.813	129012' -0.173	92 0.0	98.233	1.767	0.1	2
QCS# 272 3.405	51058' 0.645	12900' 0.521	120 0.0	90.435	9.565	0.1	1	QCS# 313 3.397	51047' 0.499	129014' 0.639	110 0.0	97.600	2.400	0.1	1
QCS# 273 -2.400	5202' 3.188	12903' 0.420	155 69.393	26.164	4.443	0.4	8	QCS# 314 2.547	51049' 0.483	129015' 0.119	122 0.0	94.622	5.378	0.2	1
QCS# 274 -0.794	5205' 2.929	12907' -0.042	135 44.193	53.323	2.484	0.2	14	QCS# 315 6.610	51052' 2.851	129018' 0.305	177 0.0	99.032	0.968	0.1	0
QCS# 275 4.929	5209 2.520	129010' 0.851	165 0.0	53.396	46.603	0.7	6	QCS# 317 7.267	51058' 3.134	129021' 0.125	200 0.0	13.956	86.044	2.4	5
QCS# 276 2.308	52012' 3.197	129014' 0.256	162 17.565	59.660	22.775	0.8	8	QCS# 318	5201' 3.134	129024' 0.125	200 0.0	9.625	90.375	2.4	6

QCS# 319	5204'	129029'	196	0.0	7.732	92.268	2.5	7	QCS# 350	51059'	130025'	430	0.0	59.420	40.580	0.6	4
7.210	2.857	0.132	0.0						4.395	1.885	1.372	0.0					
QCS# 320	51059'	129041'	183	0.0	99.853	0.147	0.0	2	QCS# 351	51058'	130032'	420	0.0	61.721	38.279	0.1	6
1.755	0.492	0.112	0.0						4.389	1.909	1.336	0.0					
QCS# 321	51056'	129039'	210	0.0	97.110	2.890	0.2	1	QCS# 352	51054'	130035'	365	57.379	40.476	2.145	0.2	3
1.933	0.788	0.874	0.0						-1.870	3.439	0.273	0.0					
QCS# 322	51052'	129036'	250	11.138	67.897	20.965	0.8	4	QCS# 353	51052'	130039'	320	86.225	17.822	1.953	0.3	4
2.666	3.034	0.425	11.138						-3.335	2.874	0.917	0.0					
QCS# 323	51049'	129033'	260	0.0	55.562	44.438	0.6	3	QCS# 354	51051'	130047'	230	99.588	0.412	0.0	0.0	0
5.354	3.075	0.563	0.0						-5.141	0.718	1.913	0.0					
QCS# 324	51045'	129030'	137	0.0	99.754	0.246	0.0	2	QCS# 356	51045'	130047'	244	(trace of gravel)				
2.109	0.681	-0.424	0.0									585	38.669	61.110	0.221	0.0	0
QCS# 325	51043'	129029'	92	79.835	20.104	0.061	0.0	4	QCS# 357	51047'	130042'	304	86.334	13.608	0.058	0.0	1
-2.460	1.915	0.398	79.835						-0.113	2.041	-0.123	0.0					
QCS# 326	51040'	129027'	73	51.470	48.455	0.076	0.0	5	QCS# 358	51049'	130038'	304	86.334	13.608	0.058	0.0	1
-1.010	2.024	0.016	51.470						-3.757	2.197	0.772	0.0					
QCS# 327	51035'	129025'	57.806	42.101	0.093	0.0	26	QCS# 359	51051'	130036'	276	(trace of gravel)					
-1.238	1.645	0.101	57.806									356	5.218	92.576	2.205	0.1	3
QCS# 328	51032'	129023'	70	27.674	72.326	0.0	0.0	3	QCS# 360	51053'	130029'	356	5.218	92.576	2.205	0.1	3
-0.481	0.899	0.718	27.674						1.395	1.517	-0.227	0.0					
QCS# 329	51027'	129022'	91	91.031	8.793	0.177	0.0	0	QCS# 361	51048'	130028'	356	100.00	0.0	0.0	0.0	0
-3.120	1.677	0.882	91.031						-6.346	0.361	0.159	0.0					
QCS# 330	51053'	130049'	212	80.146	19.609	0.245	0.0	0	QCS# 362	51044'	130025'	530	0.0	93.820	6.180	0.2	3
-2.702	2.569	0.652	80.146						1.595	1.568	0.151	0.0					
QCS# 331	51056'	130043'	265	31.322	67.125	1.552	0.1	3	QCS# 363	51040'	130022'	254	(trace of gravel)				
0.207	2.874	-0.306	31.322									494	(trace of gravel)				
QCS# 332	51057'	130038'	283	35.864	62.127	2.009	0.2	2	QCS# 364	51037'	130018'	375	86.797	12.089	1.114	0.4	4
0.249	2.641	-0.223	35.864						QCS# 365	51039'	130012'	375	86.797	12.089	1.114	0.4	4
QCS# 333	5200'	130034'	228	58.343	39.977	1.679	0.0	0	QCS# 366	51043'	130016'	238	(trace of gravel)				
-1.256	2.046	0.314	58.343						-4.147	2.575	1.184	0.0					
QCS# 334	5202'	130026'	366	46.705	37.556	15.739	0.8	3	QCS# 367	51046'	130019'	242	43.492	54.425	2.083	0.0	12
-0.199	4.287	0.238	46.705						QCS# 368	51050'	130022'	230	99.963	0.037	0.0	0.0	0
QCS# 335	52014'	129046'	197	0.0	96.200	3.800	0.2	1	QCS# 369	51056'	130022'	330	0.0	92.157	7.843	0.3	4
2.950	0.563	0.829	0.0						3.145	0.540	0.849	0.0					
QCS# 336	52017'	129049'	192	0.0	99.604	0.396	0.0	1	QCS# 370	51058'	130016'	284	0.0	89.211	10.789	0.4	4
2.036	0.845	-0.129	0.0						3.573	1.072	2.634	0.0					
QCS# 338	52010'	129050'	183	0.0	89.137	10.863	0.3	2	QCS# 371	5200'	13008'	155	49.567	47.001	3.432	0.0	48
3.380	1.870	1.623	0.0						-0.757	3.156	0.034	0.0					
QCS# 339	5209'	129057'	148	0.0	92.949	7.051	0.3	7	QCS# 372	51057'	13006'	146	65.530	29.877	4.593	0.0	0
3.153	0.847	-0.382	0.0						-2.192	3.641	0.347	0.0					
QCS# 340	52013'	129059'	141	0.0	96.044	3.956	0.2	4	QCS# 373	51055'	130010'	175	11.661	46.246	42.093	0.1	22
2.908	0.728	-0.043	0.0						4.010	3.650	0.190	0.0					
QCS# 341	52010'	13006'	206	0.0	79.609	20.392	0.5	5	QCS# 374	51054'	130014'	200	52.137	41.312	6.550	0.3	16
3.937	1.705	1.598	0.0						-0.901	4.151	0.243	0.0					
QCS# 342	5207'	13004'	168	0.0	96.707	3.293	0.1	2	QCS# 375	51050'	130011'	197	30.010	54.228	15.761	0.0	0
2.407	0.832	0.365	0.0						1.168	3.808	0.375	0.0					
QCS# 343	5205'	13009'	234	0.0	95.104	4.896	0.3	2	QCS# 376	51052'	13006'	175	41.245	45.245	13.410	0.0	34
2.522	0.773	0.655	0.0						0.230	4.397	0.184	0.0					
QCS# 344	5208'	130011'	292	0.0	73.223	26.777	0.4	3	QCS# 377	51053'	13002'	159	66.911	31.738	1.351	0.1	21
4.143	1.682	1.600	0.0						-1.744	2.910	0.368	0.0					
QCS# 345	52011'	130013'	480	0.0	35.034	64.966	1.6	4	QCS# 378	51050'	129059'	174	0.0	99.402	0.598	0.1	1
6.619	3.540	0.096	0.0						2.194	0.710	-0.086	0.0					
QCS# 346	5207'	130016'	440	0.0	94.988	5.012	0.1	1	QCS# 379	51048'	13003'	190	0.0	95.914	4.086	0.2	2
2.244	0.781	0.985	0.0						2.754	0.657	0.770	0.0					
QCS# 347	5204'	130022'	370	0.0	85.987	14.013	0.3	1	QCS# 381	51043'	13009'	267	17.838	74.779	7.383	0.3	3
2.796	2.150	1.279	0.0						1.491	3.333	-0.454	0.0					
QCS# 348	5203'	130014'	310	0.0	84.354	15.646	0.3	3	QCS# 382	51044'	13002'	310	0.0	57.941	42.059	1.1	4
2.965	1.545	1.304	0.0						4.648	2.258	1.036	0.0					
QCS# 349	5201'	130019'	370	0.0	86.610	13.390	0.2	1									
2.734	1.440	1.397	0.0														

QCS# 383	51047'	129057'	270	0.0	70.409	29.591	0.5	2	QCS# 420	51033'	129016'	47	97.573	2.395	0.031	0.0	0
4.381	2.345	1.168	0.0						-3.535	1.005	1.092	183 (camera station)					
QCS# 384	51044'	129054'	344	0.0	27.486	72.515	1.2	5	QCS# 424	51013'	12904'	250 (camera station)					
5.177	2.167	0.971	0.0						QCS# 427	5105'	129028'	77 (camera station)					
QCS# 385	51041'	129051'	336	0.0	50.749	49.251	1.0	4	QCS# 429	50058'	128058'	38 (camera station)					
4.443	1.960	1.203	0.0						QCS# 430	51034'	128038'	73 (camera station)					
QCS# 386	51038'	129046'	192	60.697	32.654	6.649	0.5	16	QCS# 435	5102'	128037'	40 (camera station)					
-1.369	3.848	0.423	60.697						QCS# 437	50051'	128022'	200	0.0	88.487	11.513	0.6	3
QCS# 387	51041'	129044'	256	0.0	75.346	24.654	0.6	5	QCS# 443	51014'	128038'	195	0.0	78.402	21.598	0.6	8
3.813	2.255	1.137	0.0						2.898	2.117	1.345	0.0					
QCS# 388	51044'	129045'	310	0.0	29.722	70.279	1.3	5	QCS# 444	51016'	128046'	135	0.0	91.121	8.878	0.3	8
5.396	2.512	0.752	0.0						3.268	2.812	0.793	0.0					
QCS# 389	51050'	129048'	252	0.0	87.763	12.237	0.5	3	QCS# 446	51022'	128023'	115 (trace of gravel)					
3.534	1.846	1.651	0.0						2.848	1.764	1.420	187 (gravity core only)					
QCS# 390	51051'	129049'	150	7.475	89.690	2.835	0.0	4	QCS# 447	51026'	128020'	40	100.000	0.0	0.0	0.0	0
1.497	1.671	-0.602	7.475						QCS# 448	51030'	128023'	40	93.945	6.055	0.0	0.0	0
QCS# 391	51055'	129055'	130 (camera station)						QCS# 450	51040'	128036'	51	83.039	16.950	0.011	0.0	0
QCS# 393	5201'	13000'	131	74.828	23.418	1.754	0.0	51	-3.695	0.457	0.046	220	0.0	92.040	7.960	0.2	1
-2.363	2.711	0.552	74.828						QCS# 451	51036'	128047'	59	89.650	10.326	0.025	0.0	0
QCS# 394	5205'	129058'	147	5.531	93.915	0.553	0.2	4	-3.281	1.465	0.346	49	54.923	44.986	0.091	0.0	0
1.125	1.108	-0.224	5.531						QCS# 452	51029'	12901'	48	40.489	59.388	0.123	0.0	9
QCS# 395	5206'	129051'	119	46.915	50.624	2.461	0.1	26	-2.901	2.142	0.626	86	0.0	99.746	0.254	0.0	0
-0.447	2.283	0.169	46.915						QCS# 453	51021'	129011'	84	0.0	99.568	0.432	0.1	2
QCS# 396	5207'	129046'	183	0.0	58.579	41.422	0.9	16	2.573	0.920	0.649	102	0.0	95.189	4.811	0.1	1
4.212	1.755	1.378	0.0						QCS# 454	51028'	129018'	126 (trace of gravel)					
QCS# 398	5203'	129049'	115	97.544	2.352	0.105	0.0	0	-3.171	1.830	0.905	128	6.572	92.976	0.452	0.0	0
-4.608	1.221	2.014	97.544						QCS# 455	51031'	129012'	222	0.0	26.125	73.874	1.0	9
QCS# 399	5201'	129054'	122	83.604	15.712	0.685	0.4	43	-0.988	1.394	0.176	148	20.156	79.266	0.578	0.0	1
-3.786	2.792	0.780	83.604						QCS# 456	51032'	12904'	135	0.0	99.522	0.478	0.0	3
QCS# 400	51056'	129053'	132	20.914	78.023	1.063	0.1	12	-0.436	1.747	-0.097	183	1.070	84.800	1.127	0.4	6
0.654	2.447	-0.548	20.914						QCS# 458	51040'	12901'	220	0.0	44.031	55.969	1.0	5
QCS# 401	51052'	129047'	192	0.0	99.699	0.301	0.1	1	2.036	0.466	0.372	210	0.0	98.952	1.048	0.1	1
1.585	0.563	0.140	0.0						QCS# 459	51043'	12905'	139	100.000	0.0	0.0	0.0	0
QCS# 402	51049'	129044'	282	0.0	80.732	19.267	0.4	2	2.684	0.439	0.327	150	13.073	85.258	1.670	0.0	0
3.190	2.125	1.190	0.0						QCS# 461	51046'	12904'	357	0.0	36.436	63.564	0.6	3
QCS# 403	51046'	129040'	282	0.0	45.282	54.718	0.7	3	3.308	0.516	0.674						
5.353	2.792	0.580	0.0						QCS# 471	52015'	129053'						
QCS# 405	51040'	129035'	117	64.003	31.615	4.381	0.7	23	QCS# 473	5206'	129053'						
-1.091	3.039	0.336	64.003						0.713	0.951	-0.212						
QCS# 406	51037'	129033'	100	95.319	4.481	0.200	0.0	0	QCS# 474	5206'	129038'						
-4.287	1.579	1.466	95.319						5.238	2.403	0.844						
QCS# 410	51056'	129043'	146	54.471	44.573	0.956	0.2	12	QCS# 480	51054'	129058'						
-0.922	2.238	0.131	54.471						1.002	2.009	-0.549						
QCS# 411	5200'	129045'	113	39.083	60.379	0.538	0.1	5	QCS# 481	51058'	129051'						
-0.392	1.584	0.060	39.083						1.465	0.956	-0.241						
QCS# 412	5202'	129040'	209	0.0	73.930	26.070	0.4	6	QCS# 482	5201'	129040'						
4.014	1.858	1.480	0.0						2.645	2.104	1.160						
QCS# 413	51059'	129045'	198	0.0	71.355	28.645	0.4	4	QCS# 484	5201'	129031'						
3.995	2.027	1.184	0.0						4.751	2.108	1.076						
QCS# 414	51056'	129035'	208	0.0	6.577	93.423	2.1	6	QCS# 485	51056'	129027'						
7.413	2.883	0.119	0.0						2.443	0.535	0.461						
QCS# 416	51048'	129026'	183	74.467	24.973	0.560	0.2	11	QCS# 486	51054'	129049'						
-2.077	2.136	0.453	74.467						-4.325	0.588	0.816						
QCS# 417	51045'	129024'	84	23.614	76.319	0.067	0.0	0	QCS# 487	51051'	129055'						
0.577	1.820	-0.450	23.614						1.012	1.710	-0.623						
QCS# 418	51042'	129020'	59	19.356	80.556	0.088	0.0	3	QCS# 489	51042'	13005'						
0.538	1.423	-0.371	19.356						5.774	2.913	0.459						
QCS# 419	51038'	129018'	51	39.529	60.386	0.085	0.9	8									
-0.294	1.589	-0.041	39.529														

38

